arXiv:1507.04973v1 [math.CO] 17 Jul 2015

Infinitely Many Nonsolvable Groups Whose Cayley Graphs are Hamiltonian

Dave Witte Morris*

Department of Mathematics and Computer Science, University of Lethbridge, Lethbridge, Alberta, T1K 6R4, Canada

Abstract. We show there are infinitely many finite groups G, such that every connected Cayley graph on G has a hamiltonian cycle, and G is not solvable. Specifically, we show that if A_5 is the alternating group on five letters, and p is any prime, such that $p \equiv 1 \pmod{30}$, then every connected Cayley graph on the direct product $A_5 \times \mathbb{Z}_p$ has a hamiltonian cycle.
Keywords. Cayley graph, hamiltonian cycle, solvable group, alternating group.
2010 AMS Classification. 05C25, 05C45

1 Introduction

It has been conjectured that every connected Cayley graph on every finite group has a hamiltonian cycle (unless the graph has less than three vertices). In support of this conjecture, the literature provides numerous infinite families of finite groups G, for which it is known that every connected Cayley graph on G has a hamiltonian cycle. (See [2] and its references for more information.) However, it seems that the union of these families contains only finitely many groups that are not solvable. This note puts an end to that unsatisfactory state of affairs:

Proposition 1.1. *There are infinitely many finite groups G, such that every connected Cayley graph on G has a hamiltonian cycle, and G is not solvable.*

Since the alternating group A_5 (of order 60) is a nonabelian simple group, and is therefore not solvable, the above is an immediate consequence of the following more specific result.

Proposition 1.2. *If p is a prime, such that $p \equiv 1 \pmod{30}$, then every connected Cayley graph on the direct product $A_5 \times \mathbb{Z}_p$ has a hamiltonian cycle.*

The proof is based on a case-by-case analysis of Cayley graphs of the group A_5. Most of the hamiltonian cycles were found by computer search (using a fairly naive backtracking algorithm).

Remark 1.3. Rather than merely groups that are not solvable, it would be much more interesting to find infinitely many finite, *simple* groups G, such that every connected Cayley graph on G has a hamiltonian cycle. Regrettably, the known methods seem to be hopelessly inadequate for this problem.

*Dave.Morris@uleth.ca, http://people.uleth.ca/∼dave.morris/

2 Preliminaries

Definition 2.1. Let S be a subset of a finite group G. The *Cayley graph* of G with respect to the connection set S is the graph $\mathrm{Cay}(G;S)$ whose vertices are the elements of G, and with edges $g \text{—} gs$ and $g \text{—} gs^{-1}$, for each $g \in G$ and $s \in S$.

Notation 2.2. Suppose S is a subset of a finite group G. For $s_1, \dots, s_m \in S \cup S^{-1}$, we use $(s_i)_{i=1}^m = (s_1, \dots, s_m)$ to denote the walk in $\mathrm{Cay}(G;S)$ that visits (in order), the vertices

$$e,\ s_1,\ s_1s_2,\ s_1s_2s_3,\ \dots,\ s_1s_2\cdots s_m.$$

We use $(s_1, \dots, s_m)^k$ to denote the concatenation of k copies of the sequence $(s_i)_{i=1}^m$, and the following illustrates other notations that are often useful:

$$(a^2, b^{-3}, s_i)_{i=1}^3 = (a, a, b^{-1}, b^{-1}, b^{-1}, s_1, a, a, b^{-1}, b^{-1}, b^{-1}, s_2, a, a, b^{-1}, b^{-1}, b^{-1}, s_3).$$

Notation 2.3. We use $\overline{\ }\colon A_5 \times \mathbb{Z}_p \to A_5$ to denote the natural projection (so $\overline{(x,y)} = x$).

Our argument in Section 3 is based on the same outline as the proof in [3] of the following result.

Lemma 2.4 ([3])**.** *Every connected Cayley graph on A_5 has a hamiltonian cycle.*

The following result is the reason that the statement of Proposition 1.2 assumes $p \equiv 1 \pmod{30}$. A much weaker hypothesis would suffice in all other parts of the proof.

Corollary 2.5. *Let S be a minimal generating set of $A_5 \times \mathbb{Z}_p$, and assume $p \equiv 1 \pmod{30}$. If there exists $a \in S$, such that $\overline{S \smallsetminus \{a\}}$ generates A_5, then $\mathrm{Cay}(A_5 \times \mathbb{Z}_p; S)$ has a hamiltonian cycle.*

Proof. Since $\gcd(|A_5|, p) = 1$, the minimality of S implies that $\langle S \smallsetminus \{a\}\rangle = A_5$. (That is, every element of $S \smallsetminus \{a\}$ projects trivially to $\mathbb{Z}_p$.) From Lemma 2.4, we know there is a hamiltonian cycle $(s_i)_{i=1}^{60}$ in $\mathrm{Cay}(A_5; S \smallsetminus \{a\})$. Since, by assumption, $p-1$ is divisible by $30 = 2 \cdot 3 \cdot 5$ (and every element of A_5 has order 1, 2, 3, or 5), we know $\overline{a}^{p-1}$ is trivial. This means $a^{p-1} \in \mathbb{Z}_p$ (so a^{p-1} centralizes A_5), so it is easy to verify that $(s_{2i-1}, a^{p-1}, s_{2i}, a^{-(p-1)})_{i=1}^{30}$ is a hamiltonian cycle in $\mathrm{Cay}(A_5 \times \mathbb{Z}_p; S)$. □ (note A.1) (note A.2)

Remark 2.6. For definiteness, we point out that we write our permutations on the left, so $gs(i) = g\big(s(i)\big)$ for $g, s \in A_5$ and $i \in \{1,2,3,4,5\}$.

The remainder of this section records a few easy consequences of the following well-known, elementary observation.

Lemma 2.7 ("Factor Group Lemma" [4, §2.2])**.** *Suppose*

- *N is a cyclic, normal subgroup of G,*
- *$(s_i)_{i=1}^m$ is a hamiltonian cycle in $\mathrm{Cay}(G/N; S)$, and*
- *the voltage $\Pi(s_i)_{i=1}^m$ generates N.*

Then $(s_1, s_2, \dots, s_m)^{|N|}$ is a hamiltonian cycle in $\mathrm{Cay}(G;S)$.

Corollary 2.8 ([2, Cor. 2.11])**.** *Suppose*

- N *is a normal subgroup of* G*, such that* $|N|$ *is prime,*
- *the image of* S *in* G/N *is a minimal generating set of* G/N*,*
- *there is a hamiltonian cycle in* $\mathrm{Cay}(G/N; S)$*, and*
- $s \equiv t \pmod N$ *for some* $s, t \in S \cup S^{-1}$ *with* $s \neq t$*.*

Then there is a hamiltonian cycle in $\mathrm{Cay}(G; S)$*.*

Corollary 2.9. *Let* S *be a minimal generating set of* $A_5 \times \mathbb{Z}_p$*, such that* $\overline{S}$ *is a minimal generating set of* A_5*. If every element of* $\overline{S}$ *has order* 2*, then* $\mathrm{Cay}(A_5 \times \mathbb{Z}_p; S)$ *has a hamiltonian cycle.*

Notation 2.10. Let $C = (s_i)_{i=1}^m$ be a walk in a Cayley graph $\mathrm{Cay}(G; S)$. For $s \in S$, we use $\mathrm{wt}_C(s)$ to denote the difference between the number of occurrences of s and the number of occurrences of s^{-1} in C. (This is the *net weight* of the generator s in C.)

Lemma 2.11. *Let* $S = \{a_1, \ldots, a_k, b_1, \ldots, b_\ell\}$ *be a minimal generating set of* $A_5 \times \mathbb{Z}_p$*, such that* $\overline{S}$ *is a minimal generating set of* A_5*. Assume*

- $\ell \geq 1$*, and* $|\overline{a_i}| = 2$ *for all* i*,*
- $C_1, \ldots, C_\ell$ *are hamiltonian cycles in* $\mathrm{Cay}(A_5; \overline{S})$*, and*
- $[\mathrm{wt}_{C_i}(b_j)]$ *is the* $\ell \times \ell$ *matrix whose* (i, j) *entry is* $\mathrm{wt}_{C_i}(b_j)$*.*

If $\det[\mathrm{wt}_{C_i}(b_j)] \not\equiv 0 \pmod p$*, then* $\mathrm{Cay}(A_5 \times \mathbb{Z}_p; S)$ *has a hamiltonian cycle.*

Proof. We may assume that $a_i \in A_5$ for $1 \leq i \leq k$, for otherwise Corollary 2.8 applies with $s = a_i$ and $t = a_i^{-1}$. Write $b_i = (\overline{b_i}, v_i)$ (with $v_i \in \mathbb{Z}_p$) for $1 \leq i \leq \ell$. Since S generates $A_5 \times \mathbb{Z}_p$, the vector $[v_1, \ldots, v_\ell]$ must be nonzero in $(\mathbb{Z}_p)^\ell$. Then, since, by assumption, the matrix $[\mathrm{wt}_{C_i}(b_j)]$ is invertible over $\mathbb{Z}_p$, this implies $[\mathrm{wt}_{C_i}(b_j)][v_1, \ldots, v_\ell]^T \neq \vec{0}$ in $(\mathbb{Z}_p)^\ell$, so there is some i, such that $\sum_{j=1}^\ell wt_{C_i}(b_j)\, v_j \neq 0$ in $\mathbb{Z}_p$. This sum is precisely the voltage ΠC_i of the walk C_i, so Lemma 2.7 provides the desired hamiltonian cycle in $\mathrm{Cay}(A_5 \times \mathbb{Z}_p; S)$. □ (note A.3)

3 Proof of Proposition 1.2

Assumptions 3.1. Let S be a minimal generating set of $A_5 \times \mathbb{Z}_p$ (and, in accordance with Notation 2.3, let $\overline{S}$ be the image of S in A_5). We may assume $\overline{S}$ is a minimal generating set of A_5, for otherwise Corollary 2.5 applies. We may also assume, for every element s of S with $|\overline{s}| = 2$, that the projection of s to $\mathbb{Z}_p$ is trivial, for otherwise Corollary 2.8 applies.

Case 1. *Assume* $\#S = 2$. Write $S = \{a, b\}$.

Subcase 1.1. *Assume* $|\overline{a}| = 2$ *and* $|\overline{b}| = 3$. By applying an automorphism of A_5, we may assume $\overline{a} = (1, 2)(3, 4)$ and $\overline{b} = (2, 4, 5)$. We have the following hamiltonian cycle in $\mathrm{Cay}(A_5; \overline{S})$: (notes A.4 A.5)

$$\big((\overline{a},\overline{b}^{\,2})^3,(\overline{a},\overline{b}^{\,-2})^3,(\overline{a},\overline{b}^{\,2},\overline{a},\overline{b}^{\,-2})^2\big)^2.$$

Each left coset of $\langle\overline{b}\rangle$ appears as consecutive vertices in this cycle, so we see that (note A.6)

$$\big((a,b^{3p-1})^3,(a,b^{-(3p-1)})^3,(a,b^{3p-1},a,b^{-(3p-1)})^2\big)^2$$

passes through all of the vertices in each left coset of $\langle b\rangle$, and is therefore a hamiltonian cycle in $\mathrm{Cay}(A_5\times\mathbb{Z}_p;S)$.

Subcase 1.2. *Assume* $|\overline{a}|=2$ *and* $|\overline{b}|=5$. By applying an automorphism of G, we may assume $\overline{b}=(1,2,3,4,5)$, and that $\overline{a}$ is either $(1,2)(3,4)$, $(1,3)(2,4)$, or $(1,4)(2,3)$. (note A.7)

- For $\overline{a}=(1,2)(3,4)$, we have the hamiltonian cycle (note A.8)

$$C=\big((\overline{a},\overline{b},\overline{a},\overline{b}^{\,4})^2,\overline{a},\overline{b}^{\,2},\overline{a},\overline{b}^{\,-1},\overline{a},\overline{b}^{\,4},\overline{a},\overline{b},(\overline{a},\overline{b}^{\,2})^3,\\ \overline{a},\overline{b}^{\,-2},\overline{a},\overline{b}^{\,4},\overline{a},\overline{b}^{\,-2},(\overline{a},\overline{b})^2,\overline{a},\overline{b}^{\,-1},\overline{a},\overline{b}^{\,4},\overline{a},\overline{b}^{\,2}\big).$$

 Since $\mathrm{wt}_C(\overline{b})=29\not\equiv 0 \pmod p$, Lemma 2.11 applies.

- For $\overline{a}=(1,3)(2,4)$, we have hamiltonian cycle (note A.9)

$$C=\big(\overline{a},\overline{b}^{\,4},\overline{a},\overline{b}^{\,-1},\overline{a},\overline{b},\overline{a},\overline{b}^{\,-1},\overline{a},\overline{b}^{\,-4},\overline{a},\overline{b}^{\,-2},(\overline{a},\overline{b}^{\,-4},\overline{a},\overline{b}^{\,2})^2,\\ \overline{a},\overline{b}^{\,-4},\overline{a},\overline{b},\overline{a},\overline{b}^{\,-1},\overline{a},\overline{b}^{\,-4},\overline{a},\overline{b}^{\,-2},\overline{a},\overline{b}^{\,-4},\overline{a},\overline{b}^{\,2}\big).$$

 Since $\mathrm{wt}_C(\overline{b})=-19\not\equiv 0 \pmod p$, Lemma 2.11 applies.

- If $\overline{a}=(1,4)(2,3)$, then $\overline{a}$ normalizes $\overline{b}$, so $\langle\overline{a},\overline{b}\rangle\neq A_5$, which contradicts the fact that S is a generating set.

Subcase 1.3. *Assume* $|\overline{a}|=|\overline{b}|=3$. By applying an automorphism of G, we may assume $\overline{a}=(1,2,3)$ and $\overline{b}=(3,4,5)$. We have the hamiltonian cycle (notes A.10, A.11)

$$C_1=\big(\overline{b},\overline{a},\overline{b}^{\,2},\overline{a}^{\,2},\overline{b}^{\,-2},\overline{a}^{\,-2},\overline{b}^{\,2},\overline{a}^{\,2},\overline{b}^{\,2},\overline{a}^{\,-2},\overline{b}^{\,-2},\overline{a}^{\,-2},\overline{b}^{\,2},\overline{a}^{\,2},\overline{b}^{\,-2},\overline{a},\overline{b}^{\,-2},\\ \overline{a}^{\,-2},\overline{b}^{\,2},\overline{a},\overline{b},\overline{a}^{\,-1},\overline{b}^{\,-2},\overline{a}^{\,2},\overline{b}^{\,2},\overline{a}^{\,2},\overline{b}^{\,-2},\overline{a}^{\,-2},\overline{b}^{\,-1},\overline{a},\overline{b},\overline{a}^{\,2},\overline{b}^{\,-1},\overline{a}^{\,-2},\overline{b}^{\,-1},\overline{a}\big).$$

Note that $\mathrm{wt}_{C_1}(\overline{a})=4$ and $\mathrm{wt}_{C_1}(\overline{b})=0$. Conjugation by the permutation $(1,4)(2,5)$ interchanges $\overline{a}$ and $\overline{b}$, and therefore yields a hamiltonian cycle C_2 with $\mathrm{wt}_{C_2}(\overline{a})=0$ and $\mathrm{wt}_{C_2}(\overline{b})=4$. Since $\det\left[\begin{smallmatrix}4&0\\0&4\end{smallmatrix}\right]=16\not\equiv 0 \pmod p$, Lemma 2.11 applies.

Subcase 1.4. *Assume* $|\overline{a}|=3$ *and* $|\overline{b}|=5$. By applying an automorphism of G (and perhaps replacing $\overline{a}$ with its inverse), we may assume $\overline{b}=(1,2,3,4,5)$, and that $\overline{a}$ is either $(1,2,3)$ or $(1,2,4)$. (note A.12)

- If $\overline{a}=(1,2,3)$, then we have the hamiltonian cycles (notes A.13, A.14)

$$C_1=\big(\overline{a}^{\,2},\overline{b}^{\,-2},\overline{a}^{\,-2},\overline{b}^{\,-1},\overline{a}^{\,-1},\overline{b}^{\,3},(\overline{a}^{\,2},\overline{b})^2,\overline{a},\overline{b}^{\,-1},(\overline{a}^{\,2},\overline{b})^2,\overline{a}^{\,2},\overline{b}^{\,-2},\overline{a}^{\,2},\overline{b},\overline{a},\overline{b}^{\,-1},\\ \overline{a}^{\,-2},\overline{b},\overline{a}^{\,2},\overline{b}^{\,-1},\overline{a}^{\,-2},\overline{b},\overline{a}^{\,-1},\overline{b}^{\,-2},\overline{a}^{\,-1},\overline{b}^{\,-1},\overline{a}^{\,-2},\overline{b},\overline{a}^{\,2},(\overline{b}^{\,-1},\overline{a}^{\,-2})^2,\overline{b}\big)$$

and

$$C_2 = \big(\overline{a}^{\,2}, \overline{b}^{\,-1}, (\overline{a}^{\,2}, \overline{b})^2, \overline{a}^{\,2}, \overline{b}^{\,-1}, \overline{a}^{\,-2}, \overline{b}, \overline{a}^{\,-2}, \overline{b}^{\,-1}, \overline{a}^{\,2}, \overline{b}, \overline{a}^{\,2}, \overline{b}^{\,-1}, \overline{a}^{\,-2}, \overline{b}^{\,2},$$
$$(\overline{a}^{\,-2}, \overline{b}^{\,-1})^2, (\overline{a}^{\,-2}, \overline{b})^2, \overline{a}^{\,2}, \overline{b}, \overline{a}^{\,2}, \overline{b}^{\,-1}, \overline{a}, \overline{b}, \overline{a}^{\,-2}, \overline{b}, \overline{a}^{\,2}, (\overline{b}^{\,-1}, \overline{a}^{\,-2})^2, \overline{b}\big)$$

Then $\big[\mathrm{wt}_{C_1}(\overline{a}), \mathrm{wt}_{C_1}(\overline{b})\big] = [5, -1]$ and $\big[\mathrm{wt}_{C_2}(\overline{a}), \mathrm{wt}_{C_2}(\overline{b})\big] = [-1, 3]$. Since $\det \left[\begin{smallmatrix} 5 & -1 \\ -1 & 3 \end{smallmatrix}\right] = 14 \not\equiv 0 \pmod{p}$, Lemma 2.11 applies.

- If $\overline{a} = (1, 2, 4)$, then we have the hamiltonian cycles (notes A.15 A.16)

$$C_1 = \big(\overline{a}^{\,2}, \overline{b}^{\,-2}, \overline{a}^{\,-2}, \overline{b}^{\,-1}, \overline{a}^{\,-1}, \overline{b}, \overline{a}^{\,2}, \overline{b}^{\,-1}, \overline{a}^{\,-2}, \overline{b}, \overline{a}^{\,-2}, \overline{b}^{\,-2}, \overline{a}^{\,-2}, \overline{b}^{\,-1}, \overline{a}^{\,-1}, \overline{b}, \overline{a}^{\,2}, \overline{b}^{\,-1}, \overline{a}^{\,2}, \overline{b},$$
$$\overline{a}^{\,-2}, \overline{b}^{\,-1}, \overline{a}^{\,2}, \overline{b}, \overline{a}^{\,2}, \overline{b}^{\,-1}, \overline{a}^{\,-2}, \overline{b}, \overline{a}^{\,-2}, \overline{b}^{\,-2}, \overline{a}^{\,-2}, \overline{b}^{\,-1}, \overline{a}^{\,-1}, \overline{b}, \overline{a}^{\,2}, \overline{b}^{\,-1}, (\overline{a}^{\,-2}, \overline{b})^2\big)$$

and

$$C_2 = \big(\overline{a}^{\,2}, \overline{b}^{\,-2}, \overline{a}^{\,-2}, \overline{b}^{\,-1}, \overline{a}^{\,-1}, \overline{b}, (\overline{a}^{\,2}, \overline{b}^{\,-1})^2, \overline{a}^{\,2}, \overline{b}^{\,-2}, \overline{a}^{\,-2}, \overline{b}^{\,-1}, \overline{a}^{\,-1}, \overline{b}, \overline{a}^{\,2}, \overline{b}^{\,-1}, \overline{a}^{\,2}, \overline{b},$$
$$\overline{a}^{\,-2}, \overline{b}^{\,-1}, \overline{a}^{\,2}, \overline{b}, \overline{a}^{\,2}, \overline{b}^{\,-1}, \overline{a}^{\,-2}, \overline{b}, \overline{a}^{\,-2}, \overline{b}^{\,-2}, \overline{a}^{\,-2}, \overline{b}^{\,-1}, \overline{a}^{\,-1}, \overline{b}, \overline{a}^{\,2}, \overline{b}^{\,-1}, (\overline{a}^{\,-2}, \overline{b})^2\big)$$

Then $\big[\mathrm{wt}_{C_1}(\overline{a}), \mathrm{wt}_{C_1}(\overline{b})\big] = [-9, -5]$ and $\big[\mathrm{wt}_{C_2}(\overline{a}), \mathrm{wt}_{C_2}(\overline{b})\big] = [-1, -7]$. Since $\det \left[\begin{smallmatrix} -9 & -5 \\ -1 & -7 \end{smallmatrix}\right] = 58 \not\equiv 0 \pmod{p}$, Lemma 2.11 applies.

Subcase 1.5. *Assume* $|\overline{a}| = |\overline{b}| = 5$. By applying an automorphism of A_5, we may assume $\overline{a} = (1, 2, 3, 4, 5)$. From Sylow's Theorems, we know that A_5 has precisely six Sylow 5-subgroups. One of them is $\langle \overline{a} \rangle$, and $\langle \overline{a} \rangle$ acts transitively on the other 5. So we may assume, after conjugating by a power of $\overline{a}$, that $\langle \overline{b} \rangle = \langle (1, 2, 3, 5, 4) \rangle$. Then, by replacing b with its inverse if necessary, we may assume $\overline{b}$ is either $(1, 2, 3, 5, 4)$ or $(1, 3, 4, 2, 5)$.

- If $\overline{b} = (1, 2, 3, 5, 4)$, then we have the hamiltonian cycle (note A.17)

$$C_1 = \big(\overline{b}, \overline{a}^{\,-1}, \overline{b}, \overline{a}, \overline{b}^{\,4}, \overline{a}, \overline{b}^{\,2}, \overline{a}^{\,-1}, \overline{b}^{\,2}, \overline{a}^{\,-1}, \overline{b}^{\,-1}, \overline{a}^{\,-1}, \overline{b}, \overline{a}^{\,-1}, \overline{b}, \overline{a}, \overline{b}^{\,2}, \overline{a}^{\,-1}, \overline{b}^{\,2}, \overline{a}^{\,-1}, \overline{b}^{\,-1},$$
$$\overline{a}, \overline{b}^{\,-2}, \overline{a}^{\,-1}, \overline{b}^{\,-4}, \overline{a}^{\,-1}, \overline{b}^{\,-1}, \overline{a}, \overline{b}^{\,-4}, \overline{a}, \overline{b}^{\,-2}, \overline{a}^{\,-1}, \overline{b}^{\,-1}, \overline{a}, \overline{b}^{\,-2}, \overline{a}^{\,-1}, \overline{b}^{\,4}, \overline{a}, \overline{b}^{\,-2}, \overline{a}\big),$$

with $\big[\mathrm{wt}_{C_1}(\overline{a}), \mathrm{wt}_{C_1}(\overline{b})\big] = [-2, 0]$. Conjugating by the permutation $(4, 5)$ interchanges $\overline{a}$ and $\overline{b}$, and therefore yields a hamiltonian cycle C_2 with $\big[\mathrm{wt}_{C_2}(\overline{a}), \mathrm{wt}_{C_2}(\overline{b})\big] = [0, -2]$. Since $\det \left[\begin{smallmatrix} -2 & 0 \\ 0 & -2 \end{smallmatrix}\right] = 4 \not\equiv 0 \pmod{p}$, Lemma 2.11 applies.

- If $\overline{b} = (1, 3, 4, 2, 5)$, then we have the hamiltonian cycles (notes A.18 A.19)

$$C_1 = \big(\overline{a}^{\,4}, \overline{b}^{\,-1}, \overline{a}^{\,-4}, \overline{b}, \overline{a}^{\,4}, \overline{b}^{\,-1}, \overline{a}^{\,2}, \overline{b}, \overline{a}^{\,-2}, \overline{b}, \overline{a}^{\,-1}, \overline{b}, \overline{a}^{\,-4}, \overline{b}^{\,-1},$$
$$\overline{a}^{\,-1}, \overline{b}^{\,-1}, \overline{a}^{\,4}, \overline{b}, (\overline{a}^{\,-4}, \overline{b}^{\,-1})^2, (\overline{a}^{\,2}, \overline{b})^2, \overline{a}^{\,4}, (\overline{b}^{\,-1}, \overline{a})^2, \overline{b}\big)$$

and

$$C_2 = \big(\overline{a}^{\,4}, \overline{b}^{\,-1}, \overline{a}^{\,-4}, \overline{b}, \overline{a}^{\,4}, \overline{b}^{\,-1}, \overline{a}^{\,2}, \overline{b}, \overline{a}^{\,-1}, \overline{b}^{\,-1}, \overline{a}^{\,-4}, \overline{b}^{\,-1}, \overline{a}, \overline{b}^{\,-1},$$
$$\overline{a}^{\,-2}, \overline{b}^{\,-1}, \overline{a}^{\,4}, \overline{b}, (\overline{a}^{\,-4}, \overline{b}^{\,-1})^2, (\overline{a}^{\,2}, \overline{b})^2, \overline{a}^{\,4}, (\overline{b}^{\,-1}, \overline{a})^2, \overline{b}\big)$$

Then $\big[\mathrm{wt}_{C_1}(\overline{a}), \mathrm{wt}_{C_1}(\overline{b})\big] = [4, 0]$ and $\big[\mathrm{wt}_{C_2}(\overline{a}), \mathrm{wt}_{C_2}(\overline{b})\big] = [6, -4]$. Since $\det \left[\begin{smallmatrix} 4 & 0 \\ 6 & -4 \end{smallmatrix}\right] = -16 \not\equiv 0 \pmod{p}$, Lemma 2.11 applies.

Case 2. *Assume* $\#S \geq 3$. Since S is minimal, it is easy to see that $\#S = 3$, so we may write $S = \{a, b, c\}$ with $|\overline{a}| \leq |\overline{b}| \leq |\overline{c}|$. (note A.20)

Subcase 2.1. *Assume* $|\overline{c}| = 5$. Since $\overline{a}$ and $\overline{b}$ cannot both normalize $\langle \overline{c} \rangle$ (but every proper subgroup of A_5 whose order is divisible by 5 has order 5 or 10), we see that either $\langle \overline{a}, \overline{c} \rangle = A_5$ or $\langle \overline{b}, \overline{c} \rangle = A_5$, which contradicts the minimality of S.

Subcase 2.2. *Assume* $|\overline{a}| = |\overline{b}| = |\overline{c}| = 3$. By applying an automorphism of A_5, and perhaps replacing some generators by their inverses, we may assume (note A.21)

$$\overline{S} = \{(1,2,5), (1,3,5), (1,4,5)\} = \{\, (1,j,n) \mid 1 < j < n \,\} \quad \text{for } n = 5.$$

So [1, App. D] provides the following hamiltonian cycle in $\mathrm{Cay}(A_5; \overline{S})$: (note A.22)

$$R_1 = \Big(((\overline{a}^{\,2}, \overline{b})^2, \overline{a}^{\,2}, \overline{c})^2, \overline{a}, \overline{b}, (\overline{b}, \overline{a}^{\,2})^2, \overline{c}^{\,2}, \overline{a}^{\,2}, \overline{c}, \overline{b}, \overline{a}, \overline{b}, \overline{c}, \overline{a}, (\overline{c}, \overline{b}^{\,2})^2, (\overline{a}^{\,2}, \overline{b})^2, \overline{a}, \overline{c}^{\,2}, \overline{a}, \overline{b}^{\,2}, (\overline{a}, \overline{c}^{\,2})^2 \Big).$$

We have $[\mathrm{wt}_{R_1}(\overline{a}), \mathrm{wt}_{R_1}(\overline{b}), \mathrm{wt}_{R_1}(\overline{c})] = [29, 17, 14]$. Conjugating by $(2,3,4)$ and $(2,3,4)^2$ cyclically permutes $\{\overline{a}, \overline{b}, \overline{c}\}$, and therefore yields hamiltonian cycles R_2 and R_3, such that

$$[\mathrm{wt}_{R_2}(\overline{a}), \mathrm{wt}_{R_2}(\overline{b}), \mathrm{wt}_{R_2}(\overline{c})] = [17, 14, 29] \text{ and } [\mathrm{wt}_{R_3}(\overline{a}), \mathrm{wt}_{R_3}(\overline{b}), \mathrm{wt}_{R_3}(\overline{c})] = [14, 29, 17].$$

Since

$$\det \begin{bmatrix} 29 & 17 & 14 \\ 17 & 14 & 29 \\ 14 & 29 & 17 \end{bmatrix} = 11,340 = 2^2 \cdot 3^4 \cdot 5 \cdot 7 \not\equiv 0 \pmod{p},$$

Lemma 2.11 applies.

Subcase 2.3. *Assume* $|\overline{a}| = 2$ *and* $|\overline{b}| = |\overline{c}| = 3$. Since $\langle \overline{b}, \overline{c} \rangle$ is a proper subgroup of A_5 that is generated by two elements of order 3, it is conjugate to A_4. So we may assume $\overline{b} = (1,2,3)$ and $\overline{c} = (1,2,4)$. And then, since S is minimal, we may assume $\overline{a} = (1,2)(4,5)$ (perhaps after conjugating by an element of S_5 that interchanges $\overline{b}$ and $\overline{c}$). (note A.23)

We have the hamiltonian cycles (notes A.24, A.25)

$$\begin{aligned} C_1 = \big(&\overline{a}, \overline{c}^{\,-1}, \overline{a}, \overline{b}, \overline{a}, \overline{c}, \overline{a}, \overline{b}^{\,2}, \overline{a}, \overline{b}, \overline{c}, \overline{b}^{\,-1}, \overline{a}, \overline{b}^{\,-2}, \overline{a}, \overline{c}^{\,-1}, \overline{a}, \overline{c}^{\,-1}, \overline{b}^{\,2}, \overline{c}, \overline{b}^{\,-2}, \overline{a}, \overline{b}^{\,-2}, \overline{c}, \overline{a}, \overline{b}, \overline{c}^{\,-1}, \\ &\overline{b}^{\,-1}, \overline{a}, \overline{c}^{\,-1}, \overline{a}, \overline{b}^{\,-2}, \overline{a}, \overline{b}^{\,-1}, \overline{c}, \overline{a}, \overline{b}^{\,2}, \overline{a}, \overline{b}, \overline{c}, \overline{b}^{\,-1}, \overline{a}, \overline{c}^{\,-1}, \overline{b}^{\,2}, \overline{c}, \overline{a}, \overline{b}, \overline{c}^{\,-1}, \overline{a}, \overline{b}^{\,-1}, \overline{a}, \overline{b}\big) \end{aligned}$$

and

$$\begin{aligned} C_2 = \big(&\overline{a}, \overline{c}^{\,-1}, \overline{a}, \overline{b}, \overline{a}, \overline{c}, \overline{a}, \overline{b}^{\,2}, \overline{a}, \overline{b}, \overline{c}, \overline{b}^{\,-1}, \overline{a}, \overline{c}^{\,-1}, \overline{b}^{\,2}, \overline{c}, \overline{a}, \overline{b}, \overline{c}^{\,-1}, \overline{b}^{\,-1}, \overline{a}, \overline{b}^{\,-2}, \overline{a}, \overline{c}^{\,-1}, \overline{b}, \overline{a}, \overline{b}^{\,2}, \\ &\overline{a}, \overline{c}, \overline{a}, \overline{b}, \overline{c}^{\,-1}, \overline{b}^{\,-1}, \overline{a}, \overline{c}^{\,-1}, \overline{a}, \overline{c}^{\,-1}, \overline{b}^{\,2}, \overline{c}, \overline{b}^{\,-2}, \overline{a}, \overline{b}^{\,-2}, \overline{c}, \overline{a}, \overline{b}^{\,2}, \overline{a}, \overline{b}, \overline{c}^{\,-1}, \overline{a}, \overline{b}^{\,-1}, \overline{a}, \overline{b}\big) \end{aligned}$$

Then $\big[\mathrm{wt}_{C_1}(\overline{b}), \mathrm{wt}_{C_1}(\overline{c})\big] = [1, 0]$ and $\big[\mathrm{wt}_{C_2}(\overline{b}), \mathrm{wt}_{C_2}(\overline{c})\big] = [7, -2]$. Since $\det \left[\begin{smallmatrix} 1 & 0 \\ 7 & -2 \end{smallmatrix}\right] = -2 \not\equiv 0 \pmod{p}$, Lemma 2.11 applies.

Subcase 2.4. *Assume* $|\overline{a}| = |b| = 2$ *and* $|\overline{c}| = 3$. We may assume $\overline{c} = (1,2,3)$.

- If $\overline{a}$ interchanges 4 and 5, then we may assume $\overline{a} = (1,2)(4,5)$. And then we may assume $\overline{b}$ is either $(1,2)(3,4)$ or $(1,3)(2,4)$. (note A.26)

– If $\overline{b} = (1,2)(3,4)$, we have the hamiltonian cycle (note A.27)

$$C = \Big(\overline{a}, \overline{c}^{\,-1}, (\overline{a},\overline{b})^2, \overline{a}, \overline{c}^{\,-1}, \overline{a}, \overline{b}, \overline{c}^{\,-1}, \overline{b}, \overline{c}, (\overline{b},\overline{a})^2, \overline{c}, (\overline{a},\overline{b})^2, \overline{a}, \overline{c}, (\overline{a},\overline{b})^2,$$
$$\overline{a}, \overline{c}^{\,2}, \overline{b}, \overline{c}^{\,-1}, \overline{b}, (\overline{a},\overline{b})^2, \overline{c}^{\,-2}, \big((\overline{a},\overline{b})^2, \overline{a}, \overline{c}^{\,-1}\big)^2, \overline{c}^{\,-1}, \overline{b}, \overline{a}, \overline{c}^{\,-1}, (\overline{a},\overline{b})^2\Big),$$

with $\mathrm{wt}_C(\overline{c}) = -5 \not\equiv 0 \pmod{p}$, so Lemma 2.11 applies.

– If $\overline{b} = (1,3)(2,4)$, we have the hamiltonian cycle (note A.28)

$$C = \Big((\overline{a},\overline{b})^4, \overline{c}, (\overline{a},\overline{b})^4, \overline{a}, \overline{c}^{\,-1}, \overline{b}, \overline{a}, \overline{b}, \overline{c}^{\,-1}, (\overline{b},\overline{a})^3, \overline{c}^{\,-1},$$
$$\overline{b}, (\overline{a},\overline{b})^2, \overline{c}, \overline{b}, \overline{a}, \overline{c}^{\,-1}, (\overline{b},\overline{a})^4, \overline{b}, \overline{c}, (\overline{a},\overline{b})^4, \overline{a}, \overline{c}^{\,-1}, \overline{b}\Big),$$

with $\mathrm{wt}_C(\overline{c}) = -2 \not\equiv 0 \pmod{p}$, so Lemma 2.11 applies.

- If neither $\overline{a}$ nor $\overline{b}$ interchanges 4 and 5, then one of them must fix 4, and the other must fix 5. We may assume $\overline{a} = (1,2)(3,4)$. And then we may assume $\overline{b}$ is either $(1,2)(3,5)$ or $(1,3)(2,5)$. (note A.29)

– If $\overline{b} = (1,2)(3,5)$, we have the hamiltonian cycle (note A.30)

$$C = \Big(\overline{a}, \overline{c}^{\,-1}, \overline{a}, \overline{c}, \overline{b}, (\overline{a},\overline{b})^2, \overline{c}, (\overline{a},\overline{b})^2, (\overline{a},\overline{c})^2, \overline{a}, \overline{b}, \overline{a}, \overline{c}^{\,-2}, \overline{a}, \overline{c}^{\,-1}, \overline{b}, \overline{a},$$
$$\overline{c}, \overline{b}, \overline{a}, \overline{b}, \overline{c}^{\,2}, \big((\overline{a},\overline{b})^2, \overline{a}, \overline{c}\big)^2, (\overline{a}, \overline{c}^{\,-1})^2, (\overline{a},\overline{b})^2, \overline{a}, \overline{c}^{\,-2}, (\overline{a},\overline{b})^2\Big),$$

with $\mathrm{wt}_C(\overline{c}) = 1 \not\equiv 0 \pmod{p}$, so Lemma 2.11 applies.

– If $\overline{b} = (1,3)(2,5)$, we have the hamiltonian cycle (note A.31)

$$C = \Big(\overline{a}, \overline{b}, \overline{c}^{\,-1}, (\overline{a},\overline{b})^2, \overline{a}, \overline{c}, \overline{b}, (\overline{a},\overline{b})^4, \overline{c}, (\overline{a}, \overline{b}, \overline{a}, \overline{c}^{\,-1})^2, \overline{b}, \overline{a}, \overline{b}, \overline{c}^{\,2},$$
$$(\overline{a},\overline{b})^2, \overline{a}, \overline{c}, \big((\overline{b},\overline{a})^2, \overline{c}^{\,-1}\big)^2, (\overline{b},\overline{a})^2, \overline{b}, \overline{c}^{\,-1}, (\overline{b},\overline{a})^2, \overline{c}^{\,-1}, \overline{b}\Big),$$

with $\mathrm{wt}_C(\overline{c}) = -2 \not\equiv 0 \pmod{p}$, so Lemma 2.11 applies.

Subcase 2.5. *Assume* $|\overline{a}| = |\overline{b}| = |\overline{c}| = 2$. Since all generators are of order 2, Corollary 2.9 applies. □

References

[1] R. Gould, R. Roth, *Cayley digraphs and $(1, j, n)$-sequencings of the alternating groups A_n*, Discrete Math. 66, 91–102, 1987. MR0900932

[2] K. Kutnar, D. Marušič, D. W. Morris, J. Morris, P. Šparl, *Hamiltonian cycles in Cayley graphs whose order has few prime factors*, Ars Math. Contemp. 5, 27–71, 2012. MR2853700

[3] K. Kutnar, D. Marušič, D. W. Morris, J. Morris, P. Šparl, *Cayley graphs on A_5 are hamiltonian*, unpublished appendix to [2]. `http://arxiv.org/src/1009.5795/anc/A5.pdf`

[4] D. Witte, J. A. Gallian, *A survey: hamiltonian cycles in Cayley graphs*, Discrete Math. 51, 293–304, 1984. MR0762322

Appendix: Notes to aid the referee

Infinitely many nonsolvable groups whose Cayley graphs are hamiltonian

by Dave Witte Morris

A.1. Since $\gcd(|A_5|, p) = 1$, we have $\overline{g} \in \langle g \rangle$ for every $g \in A_5 \times \mathbb{Z}_p$. Therefore $A_5 = \overline{\langle S \smallsetminus \{a\} \rangle} \subseteq \langle S \smallsetminus \{a\} \rangle$. Since the minimality of S implies $a \notin \subseteq \langle S \smallsetminus \{a\} \rangle$, we conclude that $\langle S \smallsetminus \{a\} \rangle = A_5$.

A.2. Since $\langle S \rangle = A_5 \times \mathbb{Z}_p$, and $S \subseteq A_5$, we know that a projects nontrivially to $\mathbb{Z}_p$. Since $\gcd(|A_5|, p) = 1$, this implies $\mathbb{Z}_p \subseteq \langle a \rangle$, so every element g of $A_5 \times \mathbb{Z}_p$ can be written (uniquely) in the form $g = xa^r$ with $x \in A_5$ and $0 \leq r \leq p-1$. Since $(s_i)_{i=1}^{60}$ is a hamiltonian cycle in a Cayley graph on A_5, we have $x = s_1 s_2 \cdots s_k$, for some k with $0 \leq k < 60$. If $k = 2i - 1$ is odd, then

$$\begin{aligned} g &= xa^r \\ &= \left(\prod_{j=1}^{i-1} s_{2j-1} s_{2j} \right) s_{2i-1} a^r \\ &= \left(\prod_{j=1}^{i-1} (s_{2j-1} a^{p-1} s_{2j} a^{-(p-1)}) \right) s_{2i-1} a^r \qquad \text{(because } a^{p-1} \in \mathbb{Z}_p \text{ is in the center of } A_5 \times \mathbb{Z}_p\text{).} \end{aligned}$$

If $k = 2i$ is even, then

$$\begin{aligned} g &= xa^r \\ &= \left(\prod_{j=1}^{i-1} s_{2j-1} s_{2j} \right) s_{2i-1} s_{2i} a^r \\ &= \left(\prod_{j=1}^{i-1} s_{2j-1} a^{p-1} s_{2j} a^{-(p-1)} \right) s_{2i-1} a^{p-1} s_{2i} a^{-(p-1-r)}. \end{aligned}$$

In either case, we see that g is one of the vertices on the walk $(s_{2i-1}, a^{p-1}, s_{2i}, a^{-(p-1)})_{i=1}^{30}$. Therefore, the walk passes through all of the vertices in $\mathrm{Cay}(A_5 \times \mathbb{Z}_p; S)$.

Also, note that the walk has the correct length $(60p)$ to be a hamiltonian cycle. Finally, by using once again the fact that a^{p-1} is in the center, we see that the terminal vertex of the walk is

$$\prod_{i=1}^{30} s_{2i-1} a^{p-1} s_{2i} a^{-(p-1)} = \prod_{i=1}^{30} s_{2i-1} s_{2i} = \prod_{i=1}^{60} s_i = e,$$

because $(s_i)_{i=1}^{60}$ is a (hamiltonian) cycle, and therefore has terminal vertex e. We conclude that the walk is a hamiltonian cycle.

A.3. Write $C_i = (s_k)_{k=1}^n$, and let $\pi = \prod_{k=1}^n s_k$ be the voltage of C_i. Since C_i is a (hamiltonian) cycle in $\mathrm{Cay}(A_5; \overline{S})$, we know that $\overline{\pi}$ is trivial, so $\pi = \sum_{k=1}^n s_k^*$, where s_k^* is the projection of s_k to $\mathbb{Z}_p$. Noting that $(s^{-1})^* = -s^*$ for $s \in S$, we have

$$\begin{aligned}
\pi &= \sum_{k=1}^{n} s_k^* \\
&= \sum_{s \in S \cup S^{-1}} (\#\text{ occurrences of } s \text{ in } C_i) \cdot s^* \\
&= \sum_{s \in S} \big((\#\text{ occurrences of } s \text{ in } C_i) - (\#\text{ occurrences of } s^{-1} \text{ in } C_i) \big) \cdot s^* \\
&= \sum_{s \in S} \mathrm{wt}_{C_i}(s)\, s^* \\
&= \sum_{j=1}^{\ell} \mathrm{wt}_{C_i}(b_j)\, v_j .
\end{aligned}$$

A.4. We may assume $\overline{a} = (1,2)(3,4)$, since every element of order 2 in A_5 is conjugate to this. Then, in order for $\langle \overline{a}, \overline{b} \rangle$ to be transitive, the support of $\overline{b}$ must contain an element of each cycle of $\overline{a}$ (including the 1-cycle (5)). So we may assume $\overline{b} = (2,4,5)$ (after conjugating by $(1,2)$ and/or $(3,4)$, if necessary).

A.5. A hamiltonian cycle in $\mathrm{Cay}(A_5; \overline{a}, \overline{b})$ with $\overline{a} = (1,2)(3,4)$ and $\overline{b} = (2,4,5)$.

$$
\begin{array}{cccccccc}
 & e & \xrightarrow{\overline{a}} & (1,2)(3,4) & \xrightarrow{\overline{b}} & (1,2,3,4,5) & \xrightarrow{\overline{b}} & (1,2,5,3,4) \\
\xrightarrow{\overline{a}} & (1,5,3) & \xrightarrow{\overline{b}} & (1,5,2,4,3) & \xrightarrow{\overline{b}} & (1,5,4,2,3) & \xrightarrow{\overline{a}} & (1,3,2,5,4) \\
\xrightarrow{\overline{b}} & (1,3,2) & \xrightarrow{\overline{b}} & (1,3,2,4,5) & \xrightarrow{\overline{a}} & (1,4,2,3,5) & \xrightarrow{\overline{b}^{-1}} & (1,4,3,5,2) \\
\xrightarrow{\overline{b}^{-1}} & (1,4)(3,5) & \xrightarrow{\overline{a}} & (1,2,4,5,3) & \xrightarrow{\overline{b}^{-1}} & (1,2,3) & \xrightarrow{\overline{b}^{-1}} & (1,2,5,4,3) \\
\xrightarrow{\overline{a}} & (1,5,4) & \xrightarrow{\overline{b}^{-1}} & (1,5)(2,4) & \xrightarrow{\overline{b}^{-1}} & (1,5,2) & \xrightarrow{\overline{a}} & (2,5)(3,4) \\
\xrightarrow{\overline{b}} & (2,3,4) & \xrightarrow{\overline{b}} & (3,4,5) & \xrightarrow{\overline{a}} & (1,2)(3,5) & \xrightarrow{\overline{b}^{-1}} & (1,2,3,5,4) \\
\xrightarrow{\overline{b}^{-1}} & (1,2,4,3,5) & \xrightarrow{\overline{a}} & (1,4,5) & \xrightarrow{\overline{b}} & (1,4)(2,5) & \xrightarrow{\overline{b}} & (1,4,2) \\
\xrightarrow{\overline{a}} & (2,4,3) & \xrightarrow{\overline{b}^{-1}} & (2,5,3) & \xrightarrow{\overline{b}^{-1}} & (2,3)(4,5) & \xrightarrow{\overline{a}} & (1,3,5,4,2) \\
\xrightarrow{\overline{b}} & (1,3,5) & \xrightarrow{\overline{b}} & (1,3,5,2,4) & \xrightarrow{\overline{a}} & (1,4,5,2,3) & \xrightarrow{\overline{b}} & (1,4,2,5,3) \\
\xrightarrow{\overline{b}} & (1,4,3) & \xrightarrow{\overline{a}} & (1,2,4) & \xrightarrow{\overline{b}} & (1,2)(4,5) & \xrightarrow{\overline{b}} & (1,2,5) \\
\xrightarrow{\overline{a}} & (1,5)(3,4) & \xrightarrow{\overline{b}^{-1}} & (1,5,3,4,2) & \xrightarrow{\overline{b}^{-1}} & (1,5,2,3,4) & \xrightarrow{\overline{a}} & (1,3)(2,5) \\
\xrightarrow{\overline{b}^{-1}} & (1,3)(4,5) & \xrightarrow{\overline{b}^{-1}} & (1,3)(2,4) & \xrightarrow{\overline{a}} & (1,4)(2,3) & \xrightarrow{\overline{b}^{-1}} & (1,4,3,2,5) \\
\xrightarrow{\overline{b}^{-1}} & (1,4,5,3,2) & \xrightarrow{\overline{a}} & (2,4)(3,5) & \xrightarrow{\overline{b}} & (3,5,4) & \xrightarrow{\overline{b}} & (2,3,5) \\
\xrightarrow{\overline{a}} & (1,3,4,5,2) & \xrightarrow{\overline{b}^{-1}} & (1,3,4) & \xrightarrow{\overline{b}^{-1}} & (1,3,4,2,5) & \xrightarrow{\overline{a}} & (1,5)(2,3) \\
\xrightarrow{\overline{b}} & (1,5,3,2,4) & \xrightarrow{\overline{b}} & (1,5,4,3,2) & \xrightarrow{\overline{a}} & (2,5,4) & \xrightarrow{\overline{b}^{-1}} & (2,4,5) \\
\xrightarrow{\overline{b}^{-1}} & e & & & & & &
\end{array}
$$

A.6. For convenience, let

$$C_1 = \big((\overline{a}, \overline{b}^{\,2})^3, (\overline{a}, \overline{b}^{\,-2})^3, (\overline{a}, \overline{b}^{\,2}, \overline{a}, \overline{b}^{\,-2})^2\big)^2$$

and

$$C_2 = \big((a, b^{3p-1})^3, (a, b^{-(3p-1)})^3, (a, b^{3p-1}, a, b^{-(3p-1)})^2\big)^2.$$

Note that $\overline{b^{3p-1}} = \overline{b}^2$ (since $|\overline{b}| = 3$).

Let x be the terminal vertex of the walk C_2. From the preceding paragraph, we see that $\overline{x}$ is the terminal vertex of the hamiltonian cycle C_1, so $\overline{x}$ is trivial. The projection of x to $\mathbb{Z}_p$ is also trivial, because $\mathrm{wt}_{C_2}(b) = 0$. Therefore, the walk C_2 is closed. Since C_2 has the correct length to be a hamiltonian, we need only show that it passes through every element of $A_5 \times \mathbb{Z}_p$.

From the fact that $\overline{b^{3p-1}} = \overline{b}^2$, we see that the vertices of $\overline{C_2}$ are precisely the same elements of A_5 as the vertices of C_1; that is, the walk $\overline{C_2}$ passes through every element of A_5. Thus, given any $v \in A_5 \times \mathbb{Z}_p$, the walk C_2 visits some vertex w with $\overline{w} = \overline{v}$; that is, v and w are in the same coset of $\mathbb{Z}_p$. Since $\mathbb{Z}_p \subseteq \langle b \rangle$, this implies that v and w are in the same left coset of $\langle b \rangle$. Also, since there are never two consecutive appearances of a in C_2, and every occurrence of b is contained in a string b^{3p-1}, we know that C_2 traverses every element of any left coset of $\langle b \rangle$ that it enters. In particular, C_2 traverses every element of the left coset of w, so it passes through v.

A.7. We may assume $\overline{b} = (1, 2, 3, 4, 5)$, after conjugating by some permutation in S_5. Since $|\overline{a}| = 2$, it has a fixed point, which we may assume is 5 (after conjugating by a power of $\overline{b}$). So $|\overline{a}|$ must be either $(1, 2)(3, 4)$, $(1, 3)(2, 4)$, or $(1, 4)(2, 3)$.

A.8. A hamiltonian cycle in $\mathrm{Cay}(A_5; \overline{a}, \overline{b})$ with $\overline{a} = (1,2)(3,4)$ and $\overline{b} = (1,2,3,4,5)$.

$$
\begin{array}{cccccccc}
 & e & \xrightarrow{\overline{a}} & (1,2)(3,4) & \xrightarrow{\overline{b}} & (2,4,5) & \xrightarrow{\overline{a}} & (1,4,3,5,2) \\
\xrightarrow{\overline{b}} & (2,5,4) & \xrightarrow{\overline{b}} & (1,5)(2,3) & \xrightarrow{\overline{b}} & (1,3,4) & \xrightarrow{\overline{b}} & (1,2,4,5,3) \\
\xrightarrow{\overline{a}} & (1,4)(3,5) & \xrightarrow{\overline{b}} & (1,2,5,4,3) & \xrightarrow{\overline{a}} & (1,5,4) & \xrightarrow{\overline{b}} & (1,2,3) \\
\xrightarrow{\overline{b}} & (1,3,4,5,2) & \xrightarrow{\overline{b}} & (2,4)(3,5) & \xrightarrow{\overline{b}} & (1,4,3,2,5) & \xrightarrow{\overline{a}} & (1,5)(2,4) \\
\xrightarrow{\overline{b}} & (1,4)(2,3) & \xrightarrow{\overline{b}} & (1,3)(4,5) & \xrightarrow{\overline{a}} & (1,2,3,5,4) & \xrightarrow{\overline{b}^{-1}} & (1,4,5) \\
\xrightarrow{\overline{a}} & (1,2,4,3,5) & \xrightarrow{\overline{b}} & (1,4)(2,5) & \xrightarrow{\overline{b}} & (1,5,4,2,3) & \xrightarrow{\overline{b}} & (1,3,2) \\
\xrightarrow{\overline{b}} & (3,4,5) & \xrightarrow{\overline{a}} & (1,2)(3,5) & \xrightarrow{\overline{b}} & (2,5)(3,4) & \xrightarrow{\overline{a}} & (1,5,2) \\
\xrightarrow{\overline{b}} & (2,3,4) & \xrightarrow{\overline{b}} & (1,3,2,4,5) & \xrightarrow{\overline{a}} & (1,4,2,3,5) & \xrightarrow{\overline{b}} & (1,3,2,5,4) \\
\xrightarrow{\overline{b}} & (1,5,3) & \xrightarrow{\overline{a}} & (1,2,5,3,4) & \xrightarrow{\overline{b}} & (1,5,2,4,3) & \xrightarrow{\overline{b}} & (1,4,2) \\
\xrightarrow{\overline{a}} & (2,4,3) & \xrightarrow{\overline{b}^{-1}} & (1,5,3,4,2) & \xrightarrow{\overline{b}^{-1}} & (1,3)(2,5) & \xrightarrow{\overline{a}} & (1,5,2,3,4) \\
\xrightarrow{\overline{b}} & (1,3)(2,4) & \xrightarrow{\overline{b}} & (1,4,5,3,2) & \xrightarrow{\overline{b}} & (3,5,4) & \xrightarrow{\overline{b}} & (1,2,5) \\
\xrightarrow{\overline{a}} & (1,5)(3,4) & \xrightarrow{\overline{b}^{-1}} & (2,5,3) & \xrightarrow{\overline{b}^{-1}} & (1,3,5,4,2) & \xrightarrow{\overline{a}} & (2,3)(4,5) \\
\xrightarrow{\overline{b}} & (1,3,5) & \xrightarrow{\overline{a}} & (1,2,3,4,5) & \xrightarrow{\overline{b}} & (1,3,5,2,4) & \xrightarrow{\overline{a}} & (1,4,5,2,3) \\
\xrightarrow{\overline{b}^{-1}} & (1,2,4) & \xrightarrow{\overline{a}} & (1,4,3) & \xrightarrow{\overline{b}} & (1,2)(4,5) & \xrightarrow{\overline{b}} & (2,3,5) \\
\xrightarrow{\overline{b}} & (1,3,4,2,5) & \xrightarrow{\overline{b}} & (1,5,3,2,4) & \xrightarrow{\overline{a}} & (1,4,2,5,3) & \xrightarrow{\overline{b}} & (1,5,4,3,2) \\
\xrightarrow{\overline{b}} & e & & & & & &
\end{array}
$$

A.9. A hamiltonian cycle in $\mathrm{Cay}(A_5; \overline{a}, \overline{b})$ with $\overline{a} = (1,3)(2,4)$ and $\overline{b} = (1,2,3,4,5)$.

$$
\begin{array}{llllllll}
 & e & \xrightarrow{\overline{a}} & (1,3)(2,4) & \xrightarrow{\overline{b}} & (1,4,5,3,2) & \xrightarrow{\overline{b}} & (3,5,4) \\
\xrightarrow{\overline{b}} & (1,2,5) & \xrightarrow{\overline{b}} & (1,5,2,3,4) & \xrightarrow{\overline{a}} & (1,4,3,5,2) & \xrightarrow{\overline{b}^{-1}} & (1,2,4,5,3) \\
\xrightarrow{\overline{a}} & (2,5,3) & \xrightarrow{\overline{b}} & (1,5)(3,4) & \xrightarrow{\overline{a}} & (1,4,2,3,5) & \xrightarrow{\overline{b}^{-1}} & (2,4,5) \\
\xrightarrow{\overline{a}} & (1,3)(2,5) & \xrightarrow{\overline{b}^{-1}} & (1,2,3,5,4) & \xrightarrow{\overline{b}^{-1}} & (1,4,5) & \xrightarrow{\overline{b}^{-1}} & (2,4,3) \\
\xrightarrow{\overline{b}^{-1}} & (1,5,3,4,2) & \xrightarrow{\overline{a}} & (1,4)(3,5) & \xrightarrow{\overline{b}^{-1}} & (1,3,2,4,5) & \xrightarrow{\overline{b}^{-1}} & (2,3,4) \\
\xrightarrow{\overline{a}} & (1,4,3) & \xrightarrow{\overline{b}^{-1}} & (1,5,3,2,4) & \xrightarrow{\overline{b}^{-1}} & (1,3,4,2,5) & \xrightarrow{\overline{b}^{-1}} & (2,3,5) \\
\xrightarrow{\overline{b}^{-1}} & (1,2)(4,5) & \xrightarrow{\overline{a}} & (1,3,2,5,4) & \xrightarrow{\overline{b}} & (1,5,3) & \xrightarrow{\overline{b}} & (1,2)(3,4) \\
\xrightarrow{\overline{a}} & (1,4)(2,3) & \xrightarrow{\overline{b}^{-1}} & (1,5)(2,4) & \xrightarrow{\overline{b}^{-1}} & (2,5)(3,4) & \xrightarrow{\overline{b}^{-1}} & (1,2)(3,5) \\
\xrightarrow{\overline{b}^{-1}} & (1,3)(4,5) & \xrightarrow{\overline{a}} & (2,5,4) & \xrightarrow{\overline{b}} & (1,5)(2,3) & \xrightarrow{\overline{b}} & (1,3,4) \\
\xrightarrow{\overline{a}} & (1,4,2) & \xrightarrow{\overline{b}^{-1}} & (1,5,2,4,3) & \xrightarrow{\overline{b}^{-1}} & (1,2,5,3,4) & \xrightarrow{\overline{b}^{-1}} & (1,3,5) \\
\xrightarrow{\overline{b}^{-1}} & (2,3)(4,5) & \xrightarrow{\overline{a}} & (1,2,5,4,3) & \xrightarrow{\overline{b}} & (1,5,2) & \xrightarrow{\overline{a}} & (1,3,5,2,4) \\
\xrightarrow{\overline{b}^{-1}} & (1,2,3,4,5) & \xrightarrow{\overline{a}} & (1,4,3,2,5) & \xrightarrow{\overline{b}^{-1}} & (2,4)(3,5) & \xrightarrow{\overline{b}^{-1}} & (1,3,4,5,2) \\
\xrightarrow{\overline{b}^{-1}} & (1,2,3) & \xrightarrow{\overline{b}^{-1}} & (1,5,4) & \xrightarrow{\overline{a}} & (1,3,5,4,2) & \xrightarrow{\overline{b}^{-1}} & (1,4,5,2,3) \\
\xrightarrow{\overline{b}^{-1}} & (1,2,4) & \xrightarrow{\overline{a}} & (1,3,2) & \xrightarrow{\overline{b}^{-1}} & (1,5,4,2,3) & \xrightarrow{\overline{b}^{-1}} & (1,4)(2,5) \\
\xrightarrow{\overline{b}^{-1}} & (1,2,4,3,5) & \xrightarrow{\overline{b}^{-1}} & (3,4,5) & \xrightarrow{\overline{a}} & (1,4,2,5,3) & \xrightarrow{\overline{b}} & (1,5,4,3,2) \\
\xrightarrow{\overline{b}} & e & & & & & &
\end{array}
$$

A.10. The union of the supports of $\overline{a}$ and $\overline{b}$ must be all of $\{1,2,3,4,5\}$, since $\langle \overline{a}, \overline{b} \rangle$ is transitive. Since each support consists of only three elements, the intersection must be a single element, which we may assume is 3. Then, by renumbering, we may assume the support of $\overline{a}$ is $\{1,2,3\}$ and the support of $\overline{b}$ is $\{3,4,5\}$. Therefore, either $\overline{a}$ or $\overline{a}^{-1}$ is $(1,2,3)$, and either $\overline{b}$ or $\overline{b}^{-1}$ is $(3,4,5)$.

A.11. A hamiltonian cycle C_1 in $\mathrm{Cay}(A_5;\overline{a},\overline{b})$ with $\overline{a}=(1,2,3)$ and $\overline{b}=(3,4,5)$.

$$
\begin{array}{llllllll}
 & e & \xrightarrow{\overline{b}} & (3,4,5) & \xrightarrow{\overline{a}} & (1,2,4,5,3) & \xrightarrow{\overline{b}} & (1,2,4,3,5) \\
\xrightarrow{\overline{b}} & (1,2,4) & \xrightarrow{\overline{a}} & (1,4)(2,3) & \xrightarrow{\overline{a}} & (1,3,4) & \xrightarrow{\overline{b}^{-1}} & (1,3,5) \\
\xrightarrow{\overline{b}^{-1}} & (1,3)(4,5) & \xrightarrow{\overline{a}^{-1}} & (2,3)(4,5) & \xrightarrow{\overline{a}^{-1}} & (1,2)(4,5) & \xrightarrow{\overline{b}} & (1,2)(3,5) \\
\xrightarrow{\overline{b}} & (1,2)(3,4) & \xrightarrow{\overline{a}} & (2,4,3) & \xrightarrow{\overline{a}} & (1,4,3) & \xrightarrow{\overline{b}} & (1,4,5) \\
\xrightarrow{\overline{b}} & (1,4)(3,5) & \xrightarrow{\overline{a}^{-1}} & (1,5,3,2,4) & \xrightarrow{\overline{a}^{-1}} & (1,2,5,3,4) & \xrightarrow{\overline{b}^{-1}} & (1,2,5) \\
\xrightarrow{\overline{b}^{-1}} & (1,2,5,4,3) & \xrightarrow{\overline{a}^{-1}} & (3,5,4) & \xrightarrow{\overline{a}^{-1}} & (1,5,4,3,2) & \xrightarrow{\overline{b}} & (1,5,2) \\
\xrightarrow{\overline{b}} & (1,5,3,4,2) & \xrightarrow{\overline{a}} & (2,4)(3,5) & \xrightarrow{\overline{a}} & (1,4,2,5,3) & \xrightarrow{\overline{b}^{-1}} & (1,4)(2,5) \\
\xrightarrow{\overline{b}^{-1}} & (1,4,3,2,5) & \xrightarrow{\overline{a}} & (1,5)(3,4) & \xrightarrow{\overline{b}^{-1}} & (1,5,3) & \xrightarrow{\overline{b}^{-1}} & (1,5,4) \\
\xrightarrow{\overline{a}^{-1}} & (1,3,2,5,4) & \xrightarrow{\overline{a}^{-1}} & (1,2,3,5,4) & \xrightarrow{\overline{b}} & (1,2,3) & \xrightarrow{\overline{b}} & (1,2,3,4,5) \\
\xrightarrow{\overline{a}} & (1,3,2,4,5) & \xrightarrow{\overline{b}} & (1,3,5,2,4) & \xrightarrow{\overline{a}^{-1}} & (1,5,2,3,4) & \xrightarrow{\overline{b}^{-1}} & (1,5)(2,3) \\
\xrightarrow{\overline{b}^{-1}} & (1,5,4,2,3) & \xrightarrow{\overline{a}} & (1,3,5,4,2) & \xrightarrow{\overline{a}} & (2,5,4) & \xrightarrow{\overline{b}} & (2,5,3) \\
\xrightarrow{\overline{b}} & (2,5)(3,4) & \xrightarrow{\overline{a}} & (1,5,2,4,3) & \xrightarrow{\overline{a}} & (1,4,3,5,2) & \xrightarrow{\overline{b}^{-1}} & (1,4,5,3,2) \\
\xrightarrow{\overline{b}^{-1}} & (1,4,2) & \xrightarrow{\overline{a}^{-1}} & (1,3)(2,4) & \xrightarrow{\overline{a}^{-1}} & (2,3,4) & \xrightarrow{\overline{b}^{-1}} & (2,3,5) \\
\xrightarrow{\overline{a}} & (1,3)(2,5) & \xrightarrow{\overline{b}} & (1,3,4,2,5) & \xrightarrow{\overline{a}} & (1,5)(2,4) & \xrightarrow{\overline{a}} & (1,4,2,3,5) \\
\xrightarrow{\overline{b}^{-1}} & (1,4,5,2,3) & \xrightarrow{\overline{a}^{-1}} & (2,4,5) & \xrightarrow{\overline{a}^{-1}} & (1,3,4,5,2) & \xrightarrow{\overline{b}^{-1}} & (1,3,2) \\
\xrightarrow{\overline{a}} & e & & & & & &
\end{array}
$$

A.12. We may assume $\overline{b}=(1,2,3,4,5)$, by replacing it with a conjugate. Then the two fixed points of the 3-cycle $\overline{a}$ are either consecutive or are separated by only one element (in circular order). Thus, after conjugating by a power of $\overline{b}$, we may assume that the fixed points of $\overline{a}$ are either 4 and 5 or 3 and 5. Hence, $\overline{a}$ is either $(1,2,3)^{\pm 1}$ or $(1,2,4)^{\pm 1}$.

A.13. A hamiltonian cycle C_1 in $\mathrm{Cay}(A_5;\overline{a},\overline{b})$ with $\overline{a}=(1,2,3)$ and $\overline{b}=(1,2,3,4,5)$.

	e	$\xrightarrow{\overline{a}}$	$(1,2,3)$	$\xrightarrow{\overline{a}}$	$(1,3,2)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,5,4,2,3)$
$\xrightarrow{\overline{b}^{-1}}$	$(1,4)(2,5)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,3,5,2,4)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,5,2,3,4)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,2,5)$
$\xrightarrow{\overline{a}^{-1}}$	$(1,3,5)$	$\xrightarrow{\overline{b}}$	$(1,2,5,3,4)$	$\xrightarrow{\overline{b}}$	$(1,5,2,4,3)$	$\xrightarrow{\overline{b}}$	$(1,4,2)$
$\xrightarrow{\overline{a}}$	$(2,3,4)$	$\xrightarrow{\overline{a}}$	$(1,3)(2,4)$	$\xrightarrow{\overline{b}}$	$(1,4,5,3,2)$	$\xrightarrow{\overline{a}}$	$(3,4,5)$
$\xrightarrow{\overline{a}}$	$(1,2,4,5,3)$	$\xrightarrow{\overline{b}}$	$(1,4,3,5,2)$	$\xrightarrow{\overline{a}}$	$(2,5)(3,4)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,2)(3,5)$
$\xrightarrow{\overline{a}}$	$(2,5,3)$	$\xrightarrow{\overline{a}}$	$(1,5,3)$	$\xrightarrow{\overline{b}}$	$(1,2)(3,4)$	$\xrightarrow{\overline{a}}$	$(2,4,3)$
$\xrightarrow{\overline{a}}$	$(1,4,3)$	$\xrightarrow{\overline{b}}$	$(1,2)(4,5)$	$\xrightarrow{\overline{a}}$	$(2,3)(4,5)$	$\xrightarrow{\overline{a}}$	$(1,3)(4,5)$
$\xrightarrow{\overline{b}^{-1}}$	$(1,4)(2,3)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,5)(2,4)$	$\xrightarrow{\overline{a}}$	$(1,4,2,3,5)$	$\xrightarrow{\overline{a}}$	$(1,3,4,2,5)$
$\xrightarrow{\overline{b}}$	$(1,5,3,2,4)$	$\xrightarrow{\overline{a}}$	$(1,4)(3,5)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,3,2,4,5)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,2,3,4,5)$
$\xrightarrow{\overline{a}^{-1}}$	$(1,4,5)$	$\xrightarrow{\overline{b}}$	$(1,2,3,5,4)$	$\xrightarrow{\overline{a}}$	$(1,3,2,5,4)$	$\xrightarrow{\overline{a}}$	$(1,5,4)$
$\xrightarrow{\overline{b}^{-1}}$	$(1,4,3,2,5)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,2,4,3,5)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,5)(3,4)$	$\xrightarrow{\overline{b}}$	$(1,2,4)$
$\xrightarrow{\overline{a}^{-1}}$	$(1,3,4)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,5)(2,3)$	$\xrightarrow{\overline{b}^{-1}}$	$(2,5,4)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,3,5,4,2)$
$\xrightarrow{\overline{b}^{-1}}$	$(1,4,5,2,3)$	$\xrightarrow{\overline{a}^{-1}}$	$(2,4,5)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,3,4,5,2)$	$\xrightarrow{\overline{b}}$	$(2,4)(3,5)$
$\xrightarrow{\overline{a}}$	$(1,4,2,5,3)$	$\xrightarrow{\overline{a}}$	$(1,5,3,4,2)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,3)(2,5)$	$\xrightarrow{\overline{a}^{-1}}$	$(2,3,5)$
$\xrightarrow{\overline{a}^{-1}}$	$(1,5,2)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,2,5,4,3)$	$\xrightarrow{\overline{a}^{-1}}$	$(3,5,4)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,5,4,3,2)$
$\xrightarrow{\overline{b}}$	e						

A.14. A second hamiltonian cycle C_2 in $\mathrm{Cay}(A_5; \overline{a}, \overline{b})$ with $\overline{a} = (1,2,3)$ and $\overline{b} = (1,2,3,4,5)$.

	e	$\xrightarrow{\overline{a}}$	$(1,2,3)$	$\xrightarrow{\overline{a}}$	$(1,3,2)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,5,4,2,3)$
$\xrightarrow{\overline{a}}$	$(1,3,5,4,2)$	$\xrightarrow{\overline{a}}$	$(2,5,4)$	$\xrightarrow{\overline{b}}$	$(1,5)(2,3)$	$\xrightarrow{\overline{a}}$	$(1,3,5)$
$\xrightarrow{\overline{a}}$	$(1,2,5)$	$\xrightarrow{\overline{b}}$	$(1,5,2,3,4)$	$\xrightarrow{\overline{a}}$	$(1,3,5,2,4)$	$\xrightarrow{\overline{a}}$	$(1,4)(2,5)$
$\xrightarrow{\overline{b}^{-1}}$	$(1,2,4,3,5)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,5)(3,4)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,4,3,2,5)$	$\xrightarrow{\overline{b}}$	$(1,5,4)$
$\xrightarrow{\overline{a}^{-1}}$	$(1,3,2,5,4)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,2,3,5,4)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,4,5)$	$\xrightarrow{\overline{a}}$	$(1,2,3,4,5)$
$\xrightarrow{\overline{a}}$	$(1,3,2,4,5)$	$\xrightarrow{\overline{b}}$	$(1,4)(3,5)$	$\xrightarrow{\overline{a}}$	$(1,2,5,3,4)$	$\xrightarrow{\overline{a}}$	$(1,5,3,2,4)$
$\xrightarrow{\overline{b}^{-1}}$	$(1,3,4,2,5)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,4,2,3,5)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,5)(2,4)$	$\xrightarrow{\overline{b}}$	$(1,4)(2,3)$
$\xrightarrow{\overline{b}}$	$(1,3)(4,5)$	$\xrightarrow{\overline{a}^{-1}}$	$(2,3)(4,5)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,2)(4,5)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,4,3)$
$\xrightarrow{\overline{a}^{-1}}$	$(2,4,3)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,2)(3,4)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,5,3)$	$\xrightarrow{\overline{a}^{-1}}$	$(2,5,3)$
$\xrightarrow{\overline{a}^{-1}}$	$(1,2)(3,5)$	$\xrightarrow{\overline{b}}$	$(2,5)(3,4)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,4,3,5,2)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,5,2,4,3)$
$\xrightarrow{\overline{b}}$	$(1,4,2)$	$\xrightarrow{\overline{a}}$	$(2,3,4)$	$\xrightarrow{\overline{a}}$	$(1,3)(2,4)$	$\xrightarrow{\overline{b}}$	$(1,4,5,3,2)$
$\xrightarrow{\overline{a}}$	$(3,4,5)$	$\xrightarrow{\overline{a}}$	$(1,2,4,5,3)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,3,4)$	$\xrightarrow{\overline{a}}$	$(1,2,4)$
$\xrightarrow{\overline{b}}$	$(1,4,5,2,3)$	$\xrightarrow{\overline{a}^{-1}}$	$(2,4,5)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,3,4,5,2)$	$\xrightarrow{\overline{b}}$	$(2,4)(3,5)$
$\xrightarrow{\overline{a}}$	$(1,4,2,5,3)$	$\xrightarrow{\overline{a}}$	$(1,5,3,4,2)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,3)(2,5)$	$\xrightarrow{\overline{a}^{-1}}$	$(2,3,5)$
$\xrightarrow{\overline{a}^{-1}}$	$(1,5,2)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,2,5,4,3)$	$\xrightarrow{\overline{a}^{-1}}$	$(3,5,4)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,5,4,3,2)$
$\xrightarrow{\overline{b}}$	e						

A.15. A hamiltonian cycle C_1 in $\mathrm{Cay}(A_5;\overline{a},\overline{b})$ with $\overline{a}=(1,2,4)$ and $\overline{b}=(1,2,3,4,5)$.

$$
\begin{array}{cccccccc}
 & e & \xrightarrow{\overline{a}} & (1,2,4) & \xrightarrow{\overline{a}} & (1,4,2) & \xrightarrow{\overline{b}^{-1}} & (1,5,2,4,3) \\
\xrightarrow{\overline{b}^{-1}} & (1,2,5,3,4) & \xrightarrow{\overline{a}^{-1}} & (3,4,5) & \xrightarrow{\overline{a}^{-1}} & (1,5,3,4,2) & \xrightarrow{\overline{b}^{-1}} & (1,3)(2,5) \\
\xrightarrow{\overline{a}^{-1}} & (1,4,5,2,3) & \xrightarrow{\overline{b}} & (1,3,5,4,2) & \xrightarrow{\overline{a}} & (3,5,4) & \xrightarrow{\overline{a}} & (1,2,3,5,4) \\
\xrightarrow{\overline{b}^{-1}} & (1,4,5) & \xrightarrow{\overline{a}^{-1}} & (1,5)(2,4) & \xrightarrow{\overline{a}^{-1}} & (1,2,5) & \xrightarrow{\overline{b}} & (1,5,2,3,4) \\
\xrightarrow{\overline{a}^{-1}} & (2,5)(3,4) & \xrightarrow{\overline{a}^{-1}} & (1,3,4,5,2) & \xrightarrow{\overline{b}^{-1}} & (1,2,3) & \xrightarrow{\overline{b}^{-1}} & (1,5,4) \\
\xrightarrow{\overline{a}^{-1}} & (2,5,4) & \xrightarrow{\overline{a}^{-1}} & (1,2)(4,5) & \xrightarrow{\overline{b}^{-1}} & (1,4,3) & \xrightarrow{\overline{a}^{-1}} & (1,3)(2,4) \\
\xrightarrow{\overline{b}} & (1,4,5,3,2) & \xrightarrow{\overline{a}} & (2,5,3) & \xrightarrow{\overline{a}} & (1,5,3,2,4) & \xrightarrow{\overline{b}^{-1}} & (1,3,4,2,5) \\
\xrightarrow{\overline{a}} & (1,5)(3,4) & \xrightarrow{\overline{a}} & (1,2,3,4,5) & \xrightarrow{\overline{b}} & (1,3,5,2,4) & \xrightarrow{\overline{a}^{-1}} & (2,3,5) \\
\xrightarrow{\overline{a}^{-1}} & (1,4,3,5,2) & \xrightarrow{\overline{b}^{-1}} & (1,2,4,5,3) & \xrightarrow{\overline{a}} & (1,4,2,5,3) & \xrightarrow{\overline{a}} & (1,5,3) \\
\xrightarrow{\overline{b}} & (1,2)(3,4) & \xrightarrow{\overline{a}} & (2,3,4) & \xrightarrow{\overline{a}} & (1,3,4) & \xrightarrow{\overline{b}^{-1}} & (1,5)(2,3) \\
\xrightarrow{\overline{a}^{-1}} & (1,4,3,2,5) & \xrightarrow{\overline{a}^{-1}} & (1,3,2,4,5) & \xrightarrow{\overline{b}} & (1,4)(3,5) & \xrightarrow{\overline{a}^{-1}} & (2,4)(3,5) \\
\xrightarrow{\overline{a}^{-1}} & (1,2)(3,5) & \xrightarrow{\overline{b}^{-1}} & (1,3)(4,5) & \xrightarrow{\overline{b}^{-1}} & (1,4)(2,3) & \xrightarrow{\overline{a}^{-1}} & (2,4,3) \\
\xrightarrow{\overline{a}^{-1}} & (1,3,2) & \xrightarrow{\overline{b}^{-1}} & (1,5,4,2,3) & \xrightarrow{\overline{a}^{-1}} & (1,2,5,4,3) & \xrightarrow{\overline{b}} & (1,5,2) \\
\xrightarrow{\overline{a}} & (2,4,5) & \xrightarrow{\overline{a}} & (1,4)(2,5) & \xrightarrow{\overline{b}^{-1}} & (1,2,4,3,5) & \xrightarrow{\overline{a}^{-1}} & (1,3,5) \\
\xrightarrow{\overline{a}^{-1}} & (1,4,2,3,5) & \xrightarrow{\overline{b}} & (1,3,2,5,4) & \xrightarrow{\overline{a}^{-1}} & (2,3)(4,5) & \xrightarrow{\overline{a}^{-1}} & (1,5,4,3,2) \\
\xrightarrow{\overline{b}} & e & & & & & & \\
\end{array}
$$

A.16. A second hamiltonian cycle C_2 in $\mathrm{Cay}(A_5; \overline{a}, \overline{b})$ with $\overline{a} = (1,2,4)$ and $\overline{b} = (1,2,3,4,5)$.

$$
\begin{array}{cccccccc}
 & e & \xrightarrow{\overline{a}} & (1,2,4) & \xrightarrow{\overline{a}} & (1,4,2) & \xrightarrow{\overline{b}^{-1}} & (1,5,2,4,3) \\
\xrightarrow{\overline{b}^{-1}} & (1,2,5,3,4) & \xrightarrow{\overline{a}^{-1}} & (3,4,5) & \xrightarrow{\overline{a}^{-1}} & (1,5,3,4,2) & \xrightarrow{\overline{b}^{-1}} & (1,3)(2,5) \\
\xrightarrow{\overline{a}^{-1}} & (1,4,5,2,3) & \xrightarrow{\overline{b}} & (1,3,5,4,2) & \xrightarrow{\overline{a}} & (3,5,4) & \xrightarrow{\overline{a}} & (1,2,3,5,4) \\
\xrightarrow{\overline{b}^{-1}} & (1,4,5) & \xrightarrow{\overline{a}} & (1,2,5) & \xrightarrow{\overline{a}} & (1,5)(2,4) & \xrightarrow{\overline{b}^{-1}} & (2,5)(3,4) \\
\xrightarrow{\overline{a}} & (1,5,2,3,4) & \xrightarrow{\overline{a}} & (1,3,4,5,2) & \xrightarrow{\overline{b}^{-1}} & (1,2,3) & \xrightarrow{\overline{b}^{-1}} & (1,5,4) \\
\xrightarrow{\overline{a}^{-1}} & (2,5,4) & \xrightarrow{\overline{a}^{-1}} & (1,2)(4,5) & \xrightarrow{\overline{b}^{-1}} & (1,4,3) & \xrightarrow{\overline{a}^{-1}} & (1,3)(2,4) \\
\xrightarrow{\overline{b}} & (1,4,5,3,2) & \xrightarrow{\overline{a}} & (2,5,3) & \xrightarrow{\overline{a}} & (1,5,3,2,4) & \xrightarrow{\overline{b}^{-1}} & (1,3,4,2,5) \\
\xrightarrow{\overline{a}} & (1,5)(3,4) & \xrightarrow{\overline{a}} & (1,2,3,4,5) & \xrightarrow{\overline{b}} & (1,3,5,2,4) & \xrightarrow{\overline{a}^{-1}} & (2,3,5) \\
\xrightarrow{\overline{a}^{-1}} & (1,4,3,5,2) & \xrightarrow{\overline{b}^{-1}} & (1,2,4,5,3) & \xrightarrow{\overline{a}} & (1,4,2,5,3) & \xrightarrow{\overline{a}} & (1,5,3) \\
\xrightarrow{\overline{b}} & (1,2)(3,4) & \xrightarrow{\overline{a}} & (2,3,4) & \xrightarrow{\overline{a}} & (1,3,4) & \xrightarrow{\overline{b}^{-1}} & (1,5)(2,3) \\
\xrightarrow{\overline{a}^{-1}} & (1,4,3,2,5) & \xrightarrow{\overline{a}^{-1}} & (1,3,2,4,5) & \xrightarrow{\overline{b}} & (1,4)(3,5) & \xrightarrow{\overline{a}^{-1}} & (2,4)(3,5) \\
\xrightarrow{\overline{a}^{-1}} & (1,2)(3,5) & \xrightarrow{\overline{b}^{-1}} & (1,3)(4,5) & \xrightarrow{\overline{b}^{-1}} & (1,4)(2,3) & \xrightarrow{\overline{a}^{-1}} & (2,4,3) \\
\xrightarrow{\overline{a}^{-1}} & (1,3,2) & \xrightarrow{\overline{b}^{-1}} & (1,5,4,2,3) & \xrightarrow{\overline{a}^{-1}} & (1,2,5,4,3) & \xrightarrow{\overline{b}} & (1,5,2) \\
\xrightarrow{\overline{a}} & (2,4,5) & \xrightarrow{\overline{a}} & (1,4)(2,5) & \xrightarrow{\overline{b}^{-1}} & (1,2,4,3,5) & \xrightarrow{\overline{a}^{-1}} & (1,3,5) \\
\xrightarrow{\overline{a}^{-1}} & (1,4,2,3,5) & \xrightarrow{\overline{b}} & (1,3,2,5,4) & \xrightarrow{\overline{a}^{-1}} & (2,3)(4,5) & \xrightarrow{\overline{a}^{-1}} & (1,5,4,3,2) \\
\xrightarrow{\overline{b}} & e & & & & & & \\
\end{array}
$$

A.17. A hamiltonian cycle C_1 in $\mathrm{Cay}(A_5;\overline{a},\overline{b})$ with $\overline{a}=(1,2,3,4,5)$ and $\overline{b}=(1,2,3,5,4)$.

	e	$\xrightarrow{\overline{b}}$	$(1,2,3,5,4)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,4,5)$	$\xrightarrow{\overline{b}}$	$(1,2,3)$
$\xrightarrow{\overline{a}}$	$(1,3,4,5,2)$	$\xrightarrow{\overline{b}}$	$(2,4,3)$	$\xrightarrow{\overline{b}}$	$(1,4)(3,5)$	$\xrightarrow{\overline{b}}$	$(1,2,5)$
$\xrightarrow{\overline{b}}$	$(1,5,4,2,3)$	$\xrightarrow{\overline{a}}$	$(1,3,2)$	$\xrightarrow{\overline{b}}$	$(3,5,4)$	$\xrightarrow{\overline{b}}$	$(1,2,5,3,4)$
$\xrightarrow{\overline{a}^{-1}}$	$(1,3,5)$	$\xrightarrow{\overline{b}}$	$(1,2,5,4,3)$	$\xrightarrow{\overline{b}}$	$(1,5,3,4,2)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,3)(2,5)$
$\xrightarrow{\overline{b}^{-1}}$	$(1,4,2,3,5)$	$\xrightarrow{\overline{a}^{-1}}$	$(2,4,5)$	$\xrightarrow{\overline{b}}$	$(1,4)(2,3)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,5)(2,4)$
$\xrightarrow{\overline{b}}$	$(1,4,5,2,3)$	$\xrightarrow{\overline{a}}$	$(1,3,5,4,2)$	$\xrightarrow{\overline{b}}$	$(2,5)(3,4)$	$\xrightarrow{\overline{b}}$	$(1,5,3,2,4)$
$\xrightarrow{\overline{a}^{-1}}$	$(1,3,4,2,5)$	$\xrightarrow{\overline{b}}$	$(1,5,2,4,3)$	$\xrightarrow{\overline{b}}$	$(1,4,5,3,2)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,3)(2,4)$
$\xrightarrow{\overline{b}^{-1}}$	$(1,2,3,4,5)$	$\xrightarrow{\overline{a}}$	$(1,3,5,2,4)$	$\xrightarrow{\overline{b}^{-1}}$	$(2,3,4)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,2)(4,5)$
$\xrightarrow{\overline{a}^{-1}}$	$(1,4,3)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,3,2,4,5)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,5,2,3,4)$	$\xrightarrow{\overline{b}^{-1}}$	$(2,5,4)$
$\xrightarrow{\overline{b}^{-1}}$	$(1,2)(3,5)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,3)(4,5)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,5)(2,3)$	$\xrightarrow{\overline{a}}$	$(1,3,4)$
$\xrightarrow{\overline{b}^{-1}}$	$(2,3)(4,5)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,5,2)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,4,2,5,3)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,2,4,3,5)$
$\xrightarrow{\overline{a}}$	$(1,4)(2,5)$	$\xrightarrow{\overline{b}^{-1}}$	$(2,4)(3,5)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,2)(3,4)$	$\xrightarrow{\overline{a}^{-1}}$	$(1,5,3)$
$\xrightarrow{\overline{b}^{-1}}$	$(1,4,3,2,5)$	$\xrightarrow{\overline{a}}$	$(1,5,4)$	$\xrightarrow{\overline{b}^{-1}}$	$(2,5,3)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,4,3,5,2)$
$\xrightarrow{\overline{a}^{-1}}$	$(1,2,4,5,3)$	$\xrightarrow{\overline{b}}$	$(1,4,2)$	$\xrightarrow{\overline{b}}$	$(2,3,5)$	$\xrightarrow{\overline{b}}$	$(1,3,2,5,4)$
$\xrightarrow{\overline{b}}$	$(1,5)(3,4)$	$\xrightarrow{\overline{a}}$	$(1,2,4)$	$\xrightarrow{\overline{b}^{-1}}$	$(3,4,5)$	$\xrightarrow{\overline{b}^{-1}}$	$(1,5,4,3,2)$
$\xrightarrow{\overline{a}}$	e						

A.18. A hamiltonian cycle C_1 in $\mathrm{Cay}(A_5;\overline{a},\overline{b})$ with $\overline{a}=(1,2,3,4,5)$ and $\overline{b}=(1,3,4,2,5)$.

$$
\begin{array}{llllllll}
 & e & \xrightarrow{\overline{a}} & (1,2,3,4,5) & \xrightarrow{\overline{a}} & (1,3,5,2,4) & \xrightarrow{\overline{a}} & (1,4,2,5,3) \\
\xrightarrow{\overline{a}} & (1,5,4,3,2) & \xrightarrow{\overline{b}^{-1}} & (1,4,2,3,5) & \xrightarrow{\overline{a}^{-1}} & (2,4,5) & \xrightarrow{\overline{a}^{-1}} & (1,2)(3,4) \\
\xrightarrow{\overline{a}^{-1}} & (1,5,3) & \xrightarrow{\overline{a}^{-1}} & (1,3,2,5,4) & \xrightarrow{\overline{b}} & (1,2,4,5,3) & \xrightarrow{\overline{a}} & (1,4,3,5,2) \\
\xrightarrow{\overline{a}} & (2,5,4) & \xrightarrow{\overline{a}} & (1,5)(2,3) & \xrightarrow{\overline{a}} & (1,3,4) & \xrightarrow{\overline{b}^{-1}} & (1,5,2) \\
\xrightarrow{\overline{a}} & (2,3,4) & \xrightarrow{\overline{a}} & (1,3,2,4,5) & \xrightarrow{\overline{b}} & (1,2)(3,5) & \xrightarrow{\overline{a}^{-1}} & (1,3)(4,5) \\
\xrightarrow{\overline{a}^{-1}} & (1,4)(2,3) & \xrightarrow{\overline{b}} & (1,2,5,4,3) & \xrightarrow{\overline{a}^{-1}} & (1,4)(3,5) & \xrightarrow{\overline{b}} & (1,5,4,2,3) \\
\xrightarrow{\overline{a}^{-1}} & (1,4)(2,5) & \xrightarrow{\overline{a}^{-1}} & (1,2,4,3,5) & \xrightarrow{\overline{a}^{-1}} & (3,4,5) & \xrightarrow{\overline{a}^{-1}} & (1,3,2) \\
\xrightarrow{\overline{b}^{-1}} & (1,5)(2,4) & \xrightarrow{\overline{a}^{-1}} & (2,5)(3,4) & \xrightarrow{\overline{b}^{-1}} & (1,2,3) & \xrightarrow{\overline{a}} & (1,3,4,5,2) \\
\xrightarrow{\overline{a}} & (2,4)(3,5) & \xrightarrow{\overline{a}} & (1,4,3,2,5) & \xrightarrow{\overline{a}} & (1,5,4) & \xrightarrow{\overline{b}} & (1,3)(2,4) \\
\xrightarrow{\overline{a}^{-1}} & (1,5,2,3,4) & \xrightarrow{\overline{a}^{-1}} & (1,2,5) & \xrightarrow{\overline{a}^{-1}} & (3,5,4) & \xrightarrow{\overline{a}^{-1}} & (1,4,5,3,2) \\
\xrightarrow{\overline{b}^{-1}} & (1,3,4,2,5) & \xrightarrow{\overline{a}^{-1}} & (2,3,5) & \xrightarrow{\overline{a}^{-1}} & (1,2)(4,5) & \xrightarrow{\overline{a}^{-1}} & (1,4,3) \\
\xrightarrow{\overline{a}^{-1}} & (1,5,3,2,4) & \xrightarrow{\overline{b}^{-1}} & (1,3,5,4,2) & \xrightarrow{\overline{a}} & (2,5,3) & \xrightarrow{\overline{a}} & (1,5)(3,4) \\
\xrightarrow{\overline{b}} & (1,4,2) & \xrightarrow{\overline{a}} & (2,3)(4,5) & \xrightarrow{\overline{a}} & (1,3,5) & \xrightarrow{\overline{b}} & (1,5,3,4,2) \\
\xrightarrow{\overline{a}} & (2,4,3) & \xrightarrow{\overline{a}} & (1,4,5) & \xrightarrow{\overline{a}} & (1,2,3,5,4) & \xrightarrow{\overline{a}} & (1,3)(2,5) \\
\xrightarrow{\overline{b}^{-1}} & (1,2,4) & \xrightarrow{\overline{a}} & (1,4,5,2,3) & \xrightarrow{\overline{b}^{-1}} & (1,2,5,3,4) & \xrightarrow{\overline{a}} & (1,5,2,4,3) \\
\xrightarrow{\overline{b}} & e & & & & & & \\
\end{array}
$$

A.19. A second hamiltonian cycle C_2 in $\mathrm{Cay}(A_5;\overline{a},\overline{b})$ with $\overline{a}=(1,2,3,4,5)$ and $\overline{b}=(1,3,4,2,5)$.

$$
\begin{array}{cccccccc}
 & e & \xrightarrow{\overline{a}} & (1,2,3,4,5) & \xrightarrow{\overline{a}} & (1,3,5,2,4) & \xrightarrow{\overline{a}} & (1,4,2,5,3) \\
\xrightarrow{\overline{a}} & (1,5,4,3,2) & \xrightarrow{\overline{b}^{-1}} & (1,4,2,3,5) & \xrightarrow{\overline{a}^{-1}} & (2,4,5) & \xrightarrow{\overline{a}^{-1}} & (1,2)(3,4) \\
\xrightarrow{\overline{a}^{-1}} & (1,5,3) & \xrightarrow{\overline{a}^{-1}} & (1,3,2,5,4) & \xrightarrow{\overline{b}} & (1,2,4,5,3) & \xrightarrow{\overline{a}} & (1,4,3,5,2) \\
\xrightarrow{\overline{a}} & (2,5,4) & \xrightarrow{\overline{a}} & (1,5)(2,3) & \xrightarrow{\overline{a}} & (1,3,4) & \xrightarrow{\overline{b}^{-1}} & (1,5,2) \\
\xrightarrow{\overline{a}} & (2,3,4) & \xrightarrow{\overline{a}} & (1,3,2,4,5) & \xrightarrow{\overline{b}} & (1,2)(3,5) & \xrightarrow{\overline{a}^{-1}} & (1,3)(4,5) \\
\xrightarrow{\overline{b}^{-1}} & (1,4)(2,5) & \xrightarrow{\overline{a}^{-1}} & (1,2,4,3,5) & \xrightarrow{\overline{a}^{-1}} & (3,4,5) & \xrightarrow{\overline{a}^{-1}} & (1,3,2) \\
\xrightarrow{\overline{a}^{-1}} & (1,5,4,2,3) & \xrightarrow{\overline{b}^{-1}} & (1,4)(3,5) & \xrightarrow{\overline{a}} & (1,2,5,4,3) & \xrightarrow{\overline{b}^{-1}} & (1,4)(2,3) \\
\xrightarrow{\overline{a}^{-1}} & (1,5)(2,4) & \xrightarrow{\overline{a}^{-1}} & (2,5)(3,4) & \xrightarrow{\overline{b}^{-1}} & (1,2,3) & \xrightarrow{\overline{a}} & (1,3,4,5,2) \\
\xrightarrow{\overline{a}} & (2,4)(3,5) & \xrightarrow{\overline{a}} & (1,4,3,2,5) & \xrightarrow{\overline{a}} & (1,5,4) & \xrightarrow{\overline{b}} & (1,3)(2,4) \\
\xrightarrow{\overline{a}^{-1}} & (1,5,2,3,4) & \xrightarrow{\overline{a}^{-1}} & (1,2,5) & \xrightarrow{\overline{a}^{-1}} & (3,5,4) & \xrightarrow{\overline{a}^{-1}} & (1,4,5,3,2) \\
\xrightarrow{\overline{b}^{-1}} & (1,3,4,2,5) & \xrightarrow{\overline{a}^{-1}} & (2,3,5) & \xrightarrow{\overline{a}^{-1}} & (1,2)(4,5) & \xrightarrow{\overline{a}^{-1}} & (1,4,3) \\
\xrightarrow{\overline{a}^{-1}} & (1,5,3,2,4) & \xrightarrow{\overline{b}^{-1}} & (1,3,5,4,2) & \xrightarrow{\overline{a}} & (2,5,3) & \xrightarrow{\overline{a}} & (1,5)(3,4) \\
\xrightarrow{\overline{b}} & (1,4,2) & \xrightarrow{\overline{a}} & (2,3)(4,5) & \xrightarrow{\overline{a}} & (1,3,5) & \xrightarrow{\overline{b}} & (1,5,3,4,2) \\
\xrightarrow{\overline{a}} & (2,4,3) & \xrightarrow{\overline{a}} & (1,4,5) & \xrightarrow{\overline{a}} & (1,2,3,5,4) & \xrightarrow{\overline{a}} & (1,3)(2,5) \\
\xrightarrow{\overline{b}^{-1}} & (1,2,4) & \xrightarrow{\overline{a}} & (1,4,5,2,3) & \xrightarrow{\overline{b}^{-1}} & (1,2,5,3,4) & \xrightarrow{\overline{a}} & (1,5,2,4,3) \\
\xrightarrow{\overline{b}} & e & & & & & &
\end{array}
$$

A.20. Write $\overline{S}=\{s_1,\dots,s_r\}$ and let $H_i=\langle s_1,\dots,s_i\rangle$. Suppose $r\geq 4$. Since $\overline{S}$ is a minimal generating set, we have $H_{i-1}\subsetneq H_i$ for each i. Since $|A_5|=2^2\cdot 3\cdot 5$ is the product of only 4 primes, we must have $r=4$ and $|H_i:H_{i-1}|$ is prime for $i=1,\dots,4$. (Recall that A_5 is simple, so the natural action of A_5 on the coset space A_5/H of any proper subgroup H must be faithful. This implies $|A_5:H|>4$, because $|A_5|>|S_4|$.) Since A_4 is the only subgroup of prime index in A_5 (up to conjugacy), we may assume $H_3=A_4$. Then H_2 must be the Sylow 2-subgroup of A_4, since that is the only subgroup of prime index. So $H_3=A_4$ is generated by s_1 and s_3, contradicting the minimality of $\overline{S}$.

A.21. The minimality of $\overline{S}$ implies that $\langle \overline{a}, \overline{b} \rangle \neq A_5$, so there must be at least two elements in the intersection of the supports of $\overline{a}$ and $\overline{b}$. The supports cannot be equal, since $\overline{a} \neq \overline{b}^{\pm 1}$. Therefore, the intersection of the support consists of two points, which we may assume are 1 and 5. Then we may assume $\overline{a} = (1,2,5)$ and $\overline{b} = (1,3,5)$. Now the support of $\overline{c}$ must contain exactly two points from the support of $\overline{a}$ and exactly two points from the support of $\overline{b}$. This implies that the support of $\overline{c}$ is $\{1,4,5\}$. so $\overline{c} = (1,4,5)^{\pm 1}$.

A.22. A hamiltonian cycle R_1 in $\mathrm{Cay}(A_5; \overline{a}, \overline{b}, \overline{c})$ with $\overline{a} = (1,2,5)$, $\overline{b} = (1,3,5)$, and $\overline{c} = (1,4,5)$.

$$
\begin{array}{cccccccc}
 & e & \xrightarrow{\overline{a}} & (1,2,5) & \xrightarrow{\overline{a}} & (1,5,2) & \xrightarrow{\overline{b}} & (1,3,2) \\
\xrightarrow{\overline{a}} & (2,5,3) & \xrightarrow{\overline{a}} & (1,5)(2,3) & \xrightarrow{\overline{b}} & (1,2,3) & \xrightarrow{\overline{a}} & (1,3)(2,5) \\
\xrightarrow{\overline{a}} & (1,5,3) & \xrightarrow{\overline{c}} & (1,4,3) & \xrightarrow{\overline{a}} & (1,2,5,4,3) & \xrightarrow{\overline{a}} & (1,5,2,4,3) \\
\xrightarrow{\overline{b}} & (2,4,3) & \xrightarrow{\overline{a}} & (1,4,3,2,5) & \xrightarrow{\overline{a}} & (1,5,4,3,2) & \xrightarrow{\overline{b}} & (1,2)(3,4) \\
\xrightarrow{\overline{a}} & (2,5)(3,4) & \xrightarrow{\overline{a}} & (1,5)(3,4) & \xrightarrow{\overline{c}} & (1,3,4) & \xrightarrow{\overline{a}} & (1,2,5,3,4) \\
\xrightarrow{\overline{b}} & (1,4)(2,5) & \xrightarrow{\overline{b}} & (1,3,2,5,4) & \xrightarrow{\overline{a}} & (1,5,3,2,4) & \xrightarrow{\overline{a}} & (1,4)(2,3) \\
\xrightarrow{\overline{b}} & (1,2,3,5,4) & \xrightarrow{\overline{a}} & (1,3,5,2,4) & \xrightarrow{\overline{a}} & (1,4)(3,5) & \xrightarrow{\overline{c}} & (3,5,4) \\
\xrightarrow{\overline{c}} & (1,3,5) & \xrightarrow{\overline{a}} & (1,2)(3,5) & \xrightarrow{\overline{a}} & (2,3,5) & \xrightarrow{\overline{c}} & (1,4,2,3,5) \\
\xrightarrow{\overline{b}} & (1,5,4,2,3) & \xrightarrow{\overline{a}} & (1,3)(2,4) & \xrightarrow{\overline{b}} & (2,4)(3,5) & \xrightarrow{\overline{c}} & (1,2,4,3,5) \\
\xrightarrow{\overline{a}} & (1,4,3,5,2) & \xrightarrow{\overline{c}} & (1,3,5,4,2) & \xrightarrow{\overline{b}} & (1,5,3,4,2) & \xrightarrow{\overline{b}} & (1,4,2) \\
\xrightarrow{\overline{c}} & (1,2)(4,5) & \xrightarrow{\overline{b}} & (1,3,4,5,2) & \xrightarrow{\overline{b}} & (1,4,5,3,2) & \xrightarrow{\overline{a}} & (2,3)(4,5) \\
\xrightarrow{\overline{a}} & (1,3,2,4,5) & \xrightarrow{\overline{b}} & (1,2,4,5,3) & \xrightarrow{\overline{a}} & (1,4,5,2,3) & \xrightarrow{\overline{a}} & (1,3)(4,5) \\
\xrightarrow{\overline{b}} & (3,4,5) & \xrightarrow{\overline{a}} & (1,2,3,4,5) & \xrightarrow{\overline{c}} & (1,5,2,3,4) & \xrightarrow{\overline{c}} & (2,3,4) \\
\xrightarrow{\overline{a}} & (1,3,4,2,5) & \xrightarrow{\overline{b}} & (1,4,2,5,3) & \xrightarrow{\overline{b}} & (2,5,4) & \xrightarrow{\overline{a}} & (1,5)(2,4) \\
\xrightarrow{\overline{c}} & (1,2,4) & \xrightarrow{\overline{c}} & (2,4,5) & \xrightarrow{\overline{a}} & (1,4,5) & \xrightarrow{\overline{c}} & (1,5,4) \\
\xrightarrow{\overline{c}} & e & & & & & & \\
\end{array}
$$

A.23. By renumbering, we may assume $\overline{b} = (1, 2, 3)$. Then, since $\langle \overline{b}, \overline{c} \rangle = A_4$, we know that 4 is in the support of $\overline{c}$, and that the other two elements of the support are in $\{1, 2, 3\}$. Therefore, after conjugating by a power of $\overline{b}$, we may assume the support of $\overline{c}$ is $\{1, 2, 4\}$. So $\overline{c} = (1, 2, 4)^{\pm 1}$.

Since $\langle \overline{a}, \overline{b}, \overline{c} \rangle = A_5$ is transitive on $\{1, 2, 3, 4, 5\}$, we know that 5 is in the support of $\overline{a}$. Since $\langle \overline{a}, \overline{b} \rangle \neq A_5$, we know that the support of $\overline{b}$ does not contain precisely one element of each of the cycles of $\overline{a}$ (and similarly for the support of $\overline{c}$).

If one of the cycles of $\overline{a}$ is disjoint from the support of $\overline{b}$, then the cycle must be $(4, 5)$. The support of $\overline{c}$ contains precisely one element of this cycle, and cannot be disjoint from the other cycle, so it must contain the entire cycle. The only 2-element subset of $\{1, 2, 3\}$ contained in the support of $\overline{c}$ is $\{1, 2\}$, so $\overline{a} = (1, 2)(3, 4)$.

We may now assume that no cycle of $\overline{a}$ is disjoint from the support of $\overline{b}$ or the support of $\overline{c}$. So the cycle $(x, 5)$ in $\overline{a}$ must be either $(1, 5)$ or $(2, 5)$. We may assume it is $(1, 5)$ (after conjugating by $(1, 2)$ and replacing $\overline{b}$ and $\overline{c}$ by their inverses, if necessary). The other cycle either is $(2, 3)$ (in which case, $\langle \overline{a}, \overline{c} \rangle = A_5$), or is either $(2, 4)$ or $(3, 4)$ (in which case, $\langle \overline{a}, \overline{b} \rangle = A_5$), which contradicts the minimality of $\overline{S}$.

A.24. A hamiltonian cycle C_1 in $\mathrm{Cay}(A_5; \overline{a}, \overline{b}, \overline{c}, \overline{c})$ with $\overline{a} = (1,2)(4,5)$, $\overline{b} = (1,2,3)$, and $\overline{c} = (1,2,4)$.

$$
\begin{array}{llllllll}
 & e & \xrightarrow{\overline{a}} & (1,2)(4,5) & \xrightarrow{\overline{c}^{-1}} & (1,5,4) & \xrightarrow{\overline{a}} & (1,2,5) \\
\xrightarrow{\overline{b}} & (1,5)(2,3) & \xrightarrow{\overline{a}} & (1,3,2,5,4) & \xrightarrow{\overline{c}} & (1,5,4,3,2) & \xrightarrow{\overline{a}} & (2,5,3) \\
\xrightarrow{\overline{b}} & (1,5,3) & \xrightarrow{\overline{b}} & (1,2)(3,5) & \xrightarrow{\overline{a}} & (3,5,4) & \xrightarrow{\overline{b}} & (1,2,5,4,3) \\
\xrightarrow{\overline{c}} & (1,5,4,2,3) & \xrightarrow{\overline{b}^{-1}} & (2,5,4) & \xrightarrow{\overline{a}} & (1,5,2) & \xrightarrow{\overline{b}^{-1}} & (1,3)(2,5) \\
\xrightarrow{\overline{b}^{-1}} & (2,3,5) & \xrightarrow{\overline{a}} & (1,3,5,4,2) & \xrightarrow{\overline{c}^{-1}} & (1,2,3,5,4) & \xrightarrow{\overline{a}} & (1,3,5) \\
\xrightarrow{\overline{c}^{-1}} & (1,4,2,3,5) & \xrightarrow{\overline{b}} & (1,3,4,2,5) & \xrightarrow{\overline{b}} & (1,5)(2,4) & \xrightarrow{\overline{c}} & (1,4,5) \\
\xrightarrow{\overline{b}^{-1}} & (1,3,2,4,5) & \xrightarrow{\overline{b}^{-1}} & (1,2,3,4,5) & \xrightarrow{\overline{a}} & (1,3,4) & \xrightarrow{\overline{b}^{-1}} & (1,4)(2,3) \\
\xrightarrow{\overline{b}^{-1}} & (1,2,4) & \xrightarrow{\overline{c}} & (1,4,2) & \xrightarrow{\overline{a}} & (2,4,5) & \xrightarrow{\overline{b}} & (1,4,5,2,3) \\
\xrightarrow{\overline{c}^{-1}} & (1,5,2,4,3) & \xrightarrow{\overline{b}^{-1}} & (2,5)(3,4) & \xrightarrow{\overline{a}} & (1,5,3,4,2) & \xrightarrow{\overline{c}^{-1}} & (1,2,5,3,4) \\
\xrightarrow{\overline{a}} & (1,5)(3,4) & \xrightarrow{\overline{b}^{-1}} & (1,4,3,2,5) & \xrightarrow{\overline{b}^{-1}} & (1,2,4,3,5) & \xrightarrow{\overline{a}} & (1,4)(3,5) \\
\xrightarrow{\overline{b}^{-1}} & (1,5,3,2,4) & \xrightarrow{\overline{c}} & (1,4,5,3,2) & \xrightarrow{\overline{a}} & (2,4,3) & \xrightarrow{\overline{b}} & (1,4,3) \\
\xrightarrow{\overline{b}} & (1,2)(3,4) & \xrightarrow{\overline{a}} & (3,4,5) & \xrightarrow{\overline{b}} & (1,2,4,5,3) & \xrightarrow{\overline{c}} & (1,4,2,5,3) \\
\xrightarrow{\overline{b}^{-1}} & (2,4)(3,5) & \xrightarrow{\overline{a}} & (1,4,3,5,2) & \xrightarrow{\overline{c}^{-1}} & (1,3,5,2,4) & \xrightarrow{\overline{b}} & (1,4)(2,5) \\
\xrightarrow{\overline{b}} & (1,5,2,3,4) & \xrightarrow{\overline{c}} & (1,3,4,5,2) & \xrightarrow{\overline{a}} & (2,3,4) & \xrightarrow{\overline{b}} & (1,3)(2,4) \\
\xrightarrow{\overline{c}^{-1}} & (1,2,3) & \xrightarrow{\overline{a}} & (1,3)(4,5) & \xrightarrow{\overline{b}^{-1}} & (2,3)(4,5) & \xrightarrow{\overline{a}} & (1,3,2) \\
\xrightarrow{\overline{b}} & e & & & & & & \\
\end{array}
$$

A.25. A second hamiltonian cycle C_2 in $\mathrm{Cay}(A_5;\overline{a},\overline{b},\overline{c})$ with $\overline{a} = (1,2)(4,5)$, $\overline{b} = (1,2,3)$, and $\overline{c} = (1,2,4)$.

$$
\begin{array}{cccccccc}
 & e & \xrightarrow{\overline{a}} & (1,2)(4,5) & \xrightarrow{\overline{c}^{-1}} & (1,5,4) & \xrightarrow{\overline{a}} & (1,2,5) \\
\xrightarrow{\overline{b}} & (1,5)(2,3) & \xrightarrow{\overline{a}} & (1,3,2,5,4) & \xrightarrow{\overline{c}} & (1,5,4,3,2) & \xrightarrow{\overline{a}} & (2,5,3) \\
\xrightarrow{\overline{b}} & (1,5,3) & \xrightarrow{\overline{b}} & (1,2)(3,5) & \xrightarrow{\overline{a}} & (3,5,4) & \xrightarrow{\overline{b}} & (1,2,5,4,3) \\
\xrightarrow{\overline{c}} & (1,5,4,2,3) & \xrightarrow{\overline{b}^{-1}} & (2,5,4) & \xrightarrow{\overline{a}} & (1,5,2) & \xrightarrow{\overline{c}^{-1}} & (1,4)(2,5) \\
\xrightarrow{\overline{b}} & (1,5,2,3,4) & \xrightarrow{\overline{b}} & (1,3,5,2,4) & \xrightarrow{\overline{c}} & (1,4,3,5,2) & \xrightarrow{\overline{a}} & (2,4)(3,5) \\
\xrightarrow{\overline{b}} & (1,4,2,5,3) & \xrightarrow{\overline{c}^{-1}} & (1,2,4,5,3) & \xrightarrow{\overline{b}^{-1}} & (3,4,5) & \xrightarrow{\overline{a}} & (1,2)(3,4) \\
\xrightarrow{\overline{b}^{-1}} & (1,4,3) & \xrightarrow{\overline{b}^{-1}} & (2,4,3) & \xrightarrow{\overline{a}} & (1,4,5,3,2) & \xrightarrow{\overline{c}^{-1}} & (1,5,3,2,4) \\
\xrightarrow{\overline{b}} & (1,4)(3,5) & \xrightarrow{\overline{a}} & (1,2,4,3,5) & \xrightarrow{\overline{b}} & (1,4,3,2,5) & \xrightarrow{\overline{b}} & (1,5)(3,4) \\
\xrightarrow{\overline{a}} & (1,2,5,3,4) & \xrightarrow{\overline{c}} & (1,5,3,4,2) & \xrightarrow{\overline{a}} & (2,5)(3,4) & \xrightarrow{\overline{b}} & (1,5,2,4,3) \\
\xrightarrow{\overline{c}^{-1}} & (1,3)(2,5) & \xrightarrow{\overline{b}^{-1}} & (2,3,5) & \xrightarrow{\overline{a}} & (1,3,5,4,2) & \xrightarrow{\overline{c}^{-1}} & (1,2,3,5,4) \\
\xrightarrow{\overline{a}} & (1,3,5) & \xrightarrow{\overline{c}^{-1}} & (1,4,2,3,5) & \xrightarrow{\overline{b}} & (1,3,4,2,5) & \xrightarrow{\overline{b}} & (1,5)(2,4) \\
\xrightarrow{\overline{c}} & (1,4,5) & \xrightarrow{\overline{b}^{-1}} & (1,3,2,4,5) & \xrightarrow{\overline{b}^{-1}} & (1,2,3,4,5) & \xrightarrow{\overline{a}} & (1,3,4) \\
\xrightarrow{\overline{b}^{-1}} & (1,4)(2,3) & \xrightarrow{\overline{b}^{-1}} & (1,2,4) & \xrightarrow{\overline{c}} & (1,4,2) & \xrightarrow{\overline{a}} & (2,4,5) \\
\xrightarrow{\overline{b}} & (1,4,5,2,3) & \xrightarrow{\overline{b}} & (1,3,4,5,2) & \xrightarrow{\overline{a}} & (2,3,4) & \xrightarrow{\overline{b}} & (1,3)(2,4) \\
\xrightarrow{\overline{c}^{-1}} & (1,2,3) & \xrightarrow{\overline{a}} & (1,3)(4,5) & \xrightarrow{\overline{b}^{-1}} & (2,3)(4,5) & \xrightarrow{\overline{a}} & (1,3,2) \\
\xrightarrow{\overline{b}} & e & & & & & &
\end{array}
$$

A.26. By assumption, one of the 2-cycles in $\overline{a}$ is $(4,5)$. The other 2-cycle must be contained in $\{1,2,3\}$, so, after conjugating by a power of $\overline{c}$, we may assume $\overline{a} = (1,2)(4,5)$.

Since $\langle \overline{a},\overline{b},\overline{c}\rangle$ is transitive on $\{1,2,3,4,5\}$, the permutation $\overline{b}$ cannot have $(4,5)$ as one of its 2-cycles. So no 2-cycle in $\overline{b}$ is disjoint from the support of $\overline{c}$. So one of the 2-cycles must be contained in the support of $\overline{c}$, which is $\{1,2,3\}$. This implies that either 4 or 5 must be fixed by $\overline{b}$. We may assume it is 5 that is fixed (after conjugating by $(4,5)$ if necessary). So $\overline{b}$ is either $(1,2)(3,4)$ or $(1,3)(2,4)$ or $(2,3)(1,4)$. Since the last two are conjugate by $(1,2)$ (which centralizes $\overline{a}$ and inverts $\overline{b}$), they do not both need to be considered.

A.27. A hamiltonian cycle in $\mathrm{Cay}(A_5; \overline{a}, \overline{b}, \overline{c})$ with $\overline{a} = (1,2)(4,5)$, $\overline{b} = (1,2)(3,4)$, and $\overline{c} = (1,2,3)$.

$$
\begin{array}{llllllll}
 & e & \xrightarrow{\overline{a}} & (1,2)(4,5) & \xrightarrow{\overline{c}^{-1}} & (1,3)(4,5) & \xrightarrow{\overline{a}} & (1,2,3) \\
\xrightarrow{\overline{b}} & (1,3,4) & \xrightarrow{\overline{a}} & (1,2,3,4,5) & \xrightarrow{\overline{b}} & (1,3,5) & \xrightarrow{\overline{a}} & (1,2,3,5,4) \\
\xrightarrow{\overline{c}^{-1}} & (1,5,4) & \xrightarrow{\overline{a}} & (1,2,5) & \xrightarrow{\overline{b}} & (1,5)(3,4) & \xrightarrow{\overline{c}^{-1}} & (1,4,3,2,5) \\
\xrightarrow{\overline{b}} & (1,5)(2,4) & \xrightarrow{\overline{c}} & (1,4,2,3,5) & \xrightarrow{\overline{b}} & (1,3,2,4,5) & \xrightarrow{\overline{a}} & (1,4)(2,3) \\
\xrightarrow{\overline{b}} & (1,3)(2,4) & \xrightarrow{\overline{a}} & (1,4,5,2,3) & \xrightarrow{\overline{c}} & (1,3,4,5,2) & \xrightarrow{\overline{a}} & (2,3,4) \\
\xrightarrow{\overline{b}} & (1,3,2) & \xrightarrow{\overline{a}} & (2,3)(4,5) & \xrightarrow{\overline{b}} & (1,3,5,4,2) & \xrightarrow{\overline{a}} & (2,3,5) \\
\xrightarrow{\overline{c}} & (1,3)(2,5) & \xrightarrow{\overline{a}} & (1,5,4,2,3) & \xrightarrow{\overline{b}} & (1,3,2,5,4) & \xrightarrow{\overline{a}} & (1,5)(2,3) \\
\xrightarrow{\overline{b}} & (1,3,4,2,5) & \xrightarrow{\overline{a}} & (1,5,2,3,4) & \xrightarrow{\overline{c}} & (1,3,5,2,4) & \xrightarrow{\overline{c}} & (1,4)(2,5) \\
\xrightarrow{\overline{b}} & (1,5,2,4,3) & \xrightarrow{\overline{c}^{-1}} & (2,5)(3,4) & \xrightarrow{\overline{b}} & (1,5,2) & \xrightarrow{\overline{a}} & (2,5,4) \\
\xrightarrow{\overline{b}} & (1,5,4,3,2) & \xrightarrow{\overline{a}} & (2,5,3) & \xrightarrow{\overline{b}} & (1,5,3,4,2) & \xrightarrow{\overline{c}^{-1}} & (1,4,2,5,3) \\
\xrightarrow{\overline{c}^{-1}} & (2,4)(3,5) & \xrightarrow{\overline{a}} & (1,4,3,5,2) & \xrightarrow{\overline{b}} & (2,4,5) & \xrightarrow{\overline{a}} & (1,4,2) \\
\xrightarrow{\overline{b}} & (2,4,3) & \xrightarrow{\overline{a}} & (1,4,5,3,2) & \xrightarrow{\overline{c}^{-1}} & (1,2,4,5,3) & \xrightarrow{\overline{a}} & (1,4,3) \\
\xrightarrow{\overline{b}} & (1,2,4) & \xrightarrow{\overline{a}} & (1,4,5) & \xrightarrow{\overline{b}} & (1,2,4,3,5) & \xrightarrow{\overline{a}} & (1,4)(3,5) \\
\xrightarrow{\overline{c}^{-1}} & (1,5,3,2,4) & \xrightarrow{\overline{c}^{-1}} & (1,2,5,3,4) & \xrightarrow{\overline{b}} & (1,5,3) & \xrightarrow{\overline{a}} & (1,2,5,4,3) \\
\xrightarrow{\overline{c}^{-1}} & (3,5,4) & \xrightarrow{\overline{a}} & (1,2)(3,5) & \xrightarrow{\overline{b}} & (3,4,5) & \xrightarrow{\overline{a}} & (1,2)(3,4) \\
\xrightarrow{\overline{b}} & e & & & & & & \\
\end{array}
$$

A.28. A hamiltonian cycle in $\mathrm{Cay}(A_5; \overline{a}, \overline{b}, \overline{c})$ with $\overline{a} = (1,2)(4,5)$, $\overline{b} = (1,3)(2,4)$, and $\overline{c} = (1,2,3)$.

$$
\begin{array}{cccccccc}
 & e & \xrightarrow{\overline{a}} & (1,2)(4,5) & \xrightarrow{\overline{b}} & (1,3,2,5,4) & \xrightarrow{\overline{a}} & (1,5)(2,3) \\
\xrightarrow{\overline{b}} & (1,2,4,3,5) & \xrightarrow{\overline{a}} & (1,4)(3,5) & \xrightarrow{\overline{b}} & (1,5,3,4,2) & \xrightarrow{\overline{a}} & (2,5)(3,4) \\
\xrightarrow{\overline{b}} & (1,4,5,2,3) & \xrightarrow{\overline{c}} & (1,3,4,5,2) & \xrightarrow{\overline{a}} & (2,3,4) & \xrightarrow{\overline{b}} & (1,4,3) \\
\xrightarrow{\overline{a}} & (1,2,4,5,3) & \xrightarrow{\overline{b}} & (2,5,3) & \xrightarrow{\overline{a}} & (1,5,4,3,2) & \xrightarrow{\overline{b}} & (1,2,3,5,4) \\
\xrightarrow{\overline{a}} & (1,3,5) & \xrightarrow{\overline{b}} & (1,5)(2,4) & \xrightarrow{\overline{a}} & (1,4)(2,5) & \xrightarrow{\overline{c}^{-1}} & (1,3,5,2,4) \\
\xrightarrow{\overline{b}} & (1,5,2) & \xrightarrow{\overline{a}} & (2,5,4) & \xrightarrow{\overline{b}} & (1,3)(4,5) & \xrightarrow{\overline{c}^{-1}} & (2,3)(4,5) \\
\xrightarrow{\overline{b}} & (1,2,5,4,3) & \xrightarrow{\overline{a}} & (1,5,3) & \xrightarrow{\overline{b}} & (2,4)(3,5) & \xrightarrow{\overline{a}} & (1,4,3,5,2) \\
\xrightarrow{\overline{b}} & (1,5,2,3,4) & \xrightarrow{\overline{a}} & (1,3,4,2,5) & \xrightarrow{\overline{c}^{-1}} & (1,4,2,3,5) & \xrightarrow{\overline{b}} & (1,5)(3,4) \\
\xrightarrow{\overline{a}} & (1,2,5,3,4) & \xrightarrow{\overline{b}} & (1,4,5,3,2) & \xrightarrow{\overline{a}} & (2,4,3) & \xrightarrow{\overline{b}} & (1,2,3) \\
\xrightarrow{\overline{c}} & (1,3,2) & \xrightarrow{\overline{b}} & (1,2,4) & \xrightarrow{\overline{a}} & (1,4,5) & \xrightarrow{\overline{c}^{-1}} & (1,3,2,4,5) \\
\xrightarrow{\overline{b}} & (1,2,5) & \xrightarrow{\overline{a}} & (1,5,4) & \xrightarrow{\overline{b}} & (1,3,5,4,2) & \xrightarrow{\overline{a}} & (2,3,5) \\
\xrightarrow{\overline{b}} & (1,5,2,4,3) & \xrightarrow{\overline{a}} & (1,4,2,5,3) & \xrightarrow{\overline{b}} & (3,4,5) & \xrightarrow{\overline{a}} & (1,2)(3,4) \\
\xrightarrow{\overline{b}} & (1,4)(2,3) & \xrightarrow{\overline{c}} & (1,3,4) & \xrightarrow{\overline{a}} & (1,2,3,4,5) & \xrightarrow{\overline{b}} & (1,4,3,2,5) \\
\xrightarrow{\overline{a}} & (1,5,3,2,4) & \xrightarrow{\overline{b}} & (1,2)(3,5) & \xrightarrow{\overline{a}} & (3,5,4) & \xrightarrow{\overline{b}} & (1,5,4,2,3) \\
\xrightarrow{\overline{a}} & (1,3)(2,5) & \xrightarrow{\overline{b}} & (2,4,5) & \xrightarrow{\overline{a}} & (1,4,2) & \xrightarrow{\overline{c}^{-1}} & (1,3)(2,4) \\
\xrightarrow{\overline{b}} & e & & & & & & \\
\end{array}
$$

A.29. Since $\langle \overline{a}, \overline{c} \rangle \neq A_5$, and $\overline{a}$ does not interchange 4 and 5, we see that $\overline{a}$ must fix either 4 or 5. Similarly for $\overline{b}$. Furthermore, $\overline{a}$ and $\overline{b}$ cannot both fix 4 (or 5), since $\langle \overline{a}, \overline{b}, \overline{c} \rangle$ is transitive. So one of them fixes 4, and the other fixes 5.

We may assume it is $\overline{a}$ that fixes 5. Then we may assume $\overline{a} = (1,2)(3,4)$, after conjugating by a power of $\overline{c}$.

Then $\overline{b}$ must be either $(1,2)(3,5)$ or $(1,3)(2,5)$ or $(1,5)(2,3)$. However, the last two are conjugate by $(1,2)$ (which centralizes $\overline{a}$ and inverts $\overline{c}$), so they do not both need to be considered.

A.30. A hamiltonian cycle in $\mathrm{Cay}(A_5; \overline{a}, \overline{b}, \overline{c})$ with $\overline{a} = (1,2)(3,4)$, $\overline{b} = (1,2)(3,5)$, and $\overline{c} = (1,2,3)$.

$$
\begin{array}{llllllll}
 & e & \xrightarrow{\overline{a}} & (1,2)(3,4) & \xrightarrow{\overline{c}^{-1}} & (1,4,3) & \xrightarrow{\overline{a}} & (1,2,4) \\
\xrightarrow{\overline{c}} & (1,4)(2,3) & \xrightarrow{\overline{b}} & (1,3,5,2,4) & \xrightarrow{\overline{a}} & (1,4,5,2,3) & \xrightarrow{\overline{b}} & (1,3,2,4,5) \\
\xrightarrow{\overline{a}} & (1,4,2,3,5) & \xrightarrow{\overline{b}} & (1,3)(2,4) & \xrightarrow{\overline{c}} & (1,4,2) & \xrightarrow{\overline{a}} & (2,4,3) \\
\xrightarrow{\overline{b}} & (1,4,3,5,2) & \xrightarrow{\overline{a}} & (2,4,5) & \xrightarrow{\overline{b}} & (1,4,5,3,2) & \xrightarrow{\overline{a}} & (2,4)(3,5) \\
\xrightarrow{\overline{c}} & (1,4,2,5,3) & \xrightarrow{\overline{a}} & (1,5,3,2,4) & \xrightarrow{\overline{c}} & (1,4)(3,5) & \xrightarrow{\overline{a}} & (1,2,4,5,3) \\
\xrightarrow{\overline{b}} & (1,4,5) & \xrightarrow{\overline{a}} & (1,2,4,3,5) & \xrightarrow{\overline{c}^{-1}} & (1,5)(3,4) & \xrightarrow{\overline{c}^{-1}} & (1,4,3,2,5) \\
\xrightarrow{\overline{a}} & (1,5)(2,4) & \xrightarrow{\overline{c}^{-1}} & (1,3,4,2,5) & \xrightarrow{\overline{b}} & (1,5,4,2,3) & \xrightarrow{\overline{a}} & (1,3,2,5,4) \\
\xrightarrow{\overline{c}} & (1,5,4) & \xrightarrow{\overline{b}} & (1,2,5,3,4) & \xrightarrow{\overline{a}} & (1,5,3) & \xrightarrow{\overline{b}} & (1,2,5) \\
\xrightarrow{\overline{c}} & (1,5)(2,3) & \xrightarrow{\overline{c}} & (1,3,5) & \xrightarrow{\overline{a}} & (1,2,3,4,5) & \xrightarrow{\overline{b}} & (1,3)(4,5) \\
\xrightarrow{\overline{a}} & (1,2,3,5,4) & \xrightarrow{\overline{b}} & (1,3,4) & \xrightarrow{\overline{a}} & (1,2,3) & \xrightarrow{\overline{c}} & (1,3,2) \\
\xrightarrow{\overline{a}} & (2,3,4) & \xrightarrow{\overline{b}} & (1,3,5,4,2) & \xrightarrow{\overline{a}} & (2,3)(4,5) & \xrightarrow{\overline{b}} & (1,3,4,5,2) \\
\xrightarrow{\overline{a}} & (2,3,5) & \xrightarrow{\overline{c}} & (1,3)(2,5) & \xrightarrow{\overline{a}} & (1,5,2,3,4) & \xrightarrow{\overline{c}^{-1}} & (1,4)(2,5) \\
\xrightarrow{\overline{a}} & (1,5,2,4,3) & \xrightarrow{\overline{c}^{-1}} & (2,5)(3,4) & \xrightarrow{\overline{a}} & (1,5,2) & \xrightarrow{\overline{b}} & (2,5,3) \\
\xrightarrow{\overline{a}} & (1,5,3,4,2) & \xrightarrow{\overline{b}} & (2,5,4) & \xrightarrow{\overline{a}} & (1,5,4,3,2) & \xrightarrow{\overline{c}^{-1}} & (1,2,5,4,3) \\
\xrightarrow{\overline{c}^{-1}} & (3,5,4) & \xrightarrow{\overline{a}} & (1,2)(4,5) & \xrightarrow{\overline{b}} & (3,4,5) & \xrightarrow{\overline{a}} & (1,2)(3,5) \\
\xrightarrow{\overline{b}} & e & & & & & &
\end{array}
$$

A.31. A hamiltonian cycle in $\mathrm{Cay}(A_5; \overline{a}, \overline{b}, \overline{c})$ with $\overline{a} = (1,2)(3,4)$, $\overline{b} = (1,3)(2,5)$, and $\overline{c} = (1,2,3)$.

$$
\begin{array}{llllllll}
 & e & \xrightarrow{\overline{a}} & (1,2)(3,4) & \xrightarrow{\overline{b}} & (1,4,3,2,5) & \xrightarrow{\overline{c}^{-1}} & (1,2,4,3,5) \\
\xrightarrow{\overline{a}} & (1,4,5) & \xrightarrow{\overline{b}} & (1,3,4,5,2) & \xrightarrow{\overline{a}} & (2,3,5) & \xrightarrow{\overline{b}} & (1,5,3) \\
\xrightarrow{\overline{a}} & (1,2,5,3,4) & \xrightarrow{\overline{c}} & (1,5,3,2,4) & \xrightarrow{\overline{b}} & (1,2,3,5,4) & \xrightarrow{\overline{a}} & (1,3)(4,5) \\
\xrightarrow{\overline{b}} & (2,4,5) & \xrightarrow{\overline{a}} & (1,4,3,5,2) & \xrightarrow{\overline{b}} & (1,5)(3,4) & \xrightarrow{\overline{a}} & (1,2,5) \\
\xrightarrow{\overline{b}} & (1,3,2) & \xrightarrow{\overline{a}} & (2,3,4) & \xrightarrow{\overline{b}} & (1,4,2,5,3) & \xrightarrow{\overline{c}} & (1,5,3,4,2) \\
\xrightarrow{\overline{a}} & (2,5,3) & \xrightarrow{\overline{b}} & (1,2,3) & \xrightarrow{\overline{a}} & (1,3,4) & \xrightarrow{\overline{c}^{-1}} & (1,4)(2,3) \\
\xrightarrow{\overline{a}} & (1,3)(2,4) & \xrightarrow{\overline{b}} & (2,5,4) & \xrightarrow{\overline{a}} & (1,5,4,3,2) & \xrightarrow{\overline{c}^{-1}} & (1,2,5,4,3) \\
\xrightarrow{\overline{b}} & (2,4,3) & \xrightarrow{\overline{a}} & (1,4,2) & \xrightarrow{\overline{b}} & (1,3,4,2,5) & \xrightarrow{\overline{c}} & (1,5)(2,4) \\
\xrightarrow{\overline{c}} & (1,4,2,3,5) & \xrightarrow{\overline{a}} & (1,3,2,4,5) & \xrightarrow{\overline{b}} & (1,2)(4,5) & \xrightarrow{\overline{a}} & (3,5,4) \\
\xrightarrow{\overline{b}} & (1,5,2,4,3) & \xrightarrow{\overline{a}} & (1,4)(2,5) & \xrightarrow{\overline{c}} & (1,5,2,3,4) & \xrightarrow{\overline{b}} & (1,4)(3,5) \\
\xrightarrow{\overline{a}} & (1,2,4,5,3) & \xrightarrow{\overline{b}} & (2,3)(4,5) & \xrightarrow{\overline{a}} & (1,3,5,4,2) & \xrightarrow{\overline{c}^{-1}} & (1,5,4,2,3) \\
\xrightarrow{\overline{b}} & (2,4)(3,5) & \xrightarrow{\overline{a}} & (1,4,5,3,2) & \xrightarrow{\overline{b}} & (1,2,3,4,5) & \xrightarrow{\overline{a}} & (1,3,5) \\
\xrightarrow{\overline{c}^{-1}} & (1,5)(2,3) & \xrightarrow{\overline{b}} & (1,2)(3,5) & \xrightarrow{\overline{a}} & (3,4,5) & \xrightarrow{\overline{b}} & (1,4,5,2,3) \\
\xrightarrow{\overline{a}} & (1,3,5,2,4) & \xrightarrow{\overline{b}} & (1,5,4) & \xrightarrow{\overline{c}^{-1}} & (1,3,2,5,4) & \xrightarrow{\overline{b}} & (1,2,4) \\
\xrightarrow{\overline{a}} & (1,4,3) & \xrightarrow{\overline{b}} & (2,5)(3,4) & \xrightarrow{\overline{a}} & (1,5,2) & \xrightarrow{\overline{c}^{-1}} & (1,3)(2,5) \\
\xrightarrow{\overline{b}} & e & & & & & & \\
\end{array}
$$